



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
25.06.2003 Bulletin 2003/26

(51) Int Cl.7: **C12N 15/80, C12N 1/15**

(21) Application number: **03075731.4**

(22) Date of filing: **30.06.1994.**

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE**

(30) Priority: **23.07.1993 EP 93202195**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
94201896.1 / 0 635 574

(71) Applicant: **DSM N.V.**
6411 TE Heerlen (NL)

(72) Inventors:
• **Selten, Gerardus C. M.**
2651 HZ Berkel en Rodenrijs (NL)

• **Swinkels, Bart W.**
2611 MX Delft (NL)
• **van Gorcom, Robertus F. M.**
3768 GW Soest (NL)

(74) Representative:
Matulewicz, Emil Rudolf Antonius
DSM N.V.,
DSM Patents & Trademarks,
P.O. Box 1
2600 MA Delft (NL)

Remarks:

This application was filed on 12 - 03 - 2003 as a
divisional application to the application mentioned
under INID code 62.

(54) **Selection marker gene free recombinant strains; a method for obtaining them and the use of these strains**

(57) The present invention discloses a selection marker free system which can be used to introduce genetic modifications in bacteria, yeasts and fungi. The system can be employed to introduce or delete desired genes or DNA fragments in the genome of the indicated host species without leaving any undesired DNA i.e. the selection marker used for selection of transformants or

other DNA used for cloning. In this way strains have been developed containing only desired genes introduced at desired chromosomal sites. Similarly, desired DNA fragments have been deleted or replaced at desired sites.

Description

Technical field

5 [0001] The present invention discloses selection marker gene free recombinant strains, a method for obtaining these strains and the use of these strains. Furthermore, the method of the present invention is used for performing strain improvement.

Background of the invention

10 [0002] There is an increasing social concern about the use of recombinant DNA technology. One of the promising application areas of recombinant DNA technology is strain improvement. Starting from the early days of fermentative production processes there has been a demand for the improvement of the productivity of the strains used for production.

15 [0003] Classical strain improvement programs for industrially employed microorganisms are primarily based on random mutagenesis followed by selection. Mutagenesis methods have been described extensively; they include the use of UV light, NTG or EMS as mutagens. These methods have been described extensively for example in "Biotechnology : a comprehensive treatise in 8 vol." Volume I, Microbial fundamentals, Chapter 5b, Verlag Chemie GmbH, Weinheim, Germany.

20 [0004] Selection methods are generally developed around a suitable assay and are of major importance in the discrimination between wild type and mutant strains.

[0005] It has turned out that these classical methods are limited in their potential for improvement. Generally speaking consecutive rounds of strain improvement yield diminishing increases in yield of desired products. This is at least partially due to the random character of the mutagenesis methods employed. Apart from desired mutations these methods also give rise to mutations which are undesirable and which may negatively influence other characteristics of the strains.

25 [0006] In view of these drawbacks it can be understood that the use of recombinant DNA methods was hailed as a considerable improvement. In general, recombinant DNA methods used in strain improvement programs aim at the increased expression of desired gene products.

30 [0007] The gene products may be proteins that are of interest themselves, on the other hand it is also possible that the encoded gene products serve as regulatory proteins in the synthesis of other products.

[0008] Strains can be improved by introducing multiple copies of desired protein encoding genes into specific host organisms. However, it is also possible to increase expression levels by introducing regulatory genes.

35 [0009] Genes are introduced using vectors that serve as vehicles for introduction of the genes. Such vectors may be plasmids, cosmids or phages. The vector may be capable of expression of the genes in which case the vector generally is self-replicating. The vector may however also only be capable of integration. Another characteristic of the vector is that, when the expression product cannot be selected easily based on altered phenotypic properties, the vector is equipped with a marker that can easily be selected for.

40 [0010] Vectors have not been isolated from all known microorganisms either since no vector could be found in the organism or since available vectors from other organisms could be used with little or no modification. The same applies to selection marker genes.

45 [0011] Widespread use and the subsequent spreading of specific marker genes has recently become debatable. This is especially due to the finding that the use of antibiotics and antibiotic selection markers gives rise to an undesired spread of strains that have become antibiotic resistant. This necessitates the continued development of novel ever more potent antibiotics.

[0012] It is therefore not surprising that there is a general tendency in large scale production to use recombinant microorganisms containing no antibiotic resistance genes or more generally as little as possible of foreign DNA.

[0013] Ideally the transformed microorganism would contain only the desired gene(s), fragments thereof or modifications in the gene and as little as possible or no further remnants of the DNA used for cloning.

Summary of the invention

[0014] The present invention discloses a selection marker gene that can easily be deleted again from the recombinant host organism. The deletion of the said marker gene is based on dominant selection.

55 [0015] The marker is used in species so diverse as bacteria, filamentous fungi and yeasts.

[0016] The advantageous activity of the selection markers used herein is based on the following two step principle:

a) the gene is integrated into the genome of the host organism and recombinant cells are selected,

b) the transformed cell is grown on a substrate, which is converted by the marker gene encoded activity to a product that is lethal to the cell.

[0017] Selected cells will be recombinant and will have deleted the selection-marker gene.

5 [0018] In general terms the present invention discloses cells, that may be animal or plant cells, and microorganisms that have a modification in the genome characterized in that the alteration is introduced using the amdS gene or the cDNA derived therefrom.

[0019] An example of a selection marker gene that can be used in this way is the acetamidase gene. Preferably, this gene is obtainable from filamentous fungi, more preferably from Aspergilli, most preferably from Aspergillus nidulans.

10 [0020] The invention further shows the introduction, deletion or modification of desired heterologous or homologous genes or DNA elements in the host organisms of choice using the acetamidase (amdS) gene as a marker. Subsequently the amdS gene is deleted. Preferably, the amdS and the desired genes are introduced site-specifically.

[0021] The invention discloses a vector containing:

- 15 a) a desired DNA fragment destined for introduction into the host genome,
b) optionally a DNA sequence that enables the vector to integrate (site-specifically) into the genome of the host strain,
c) a gene encoding an acetamidase (e.g. the amdS gene from A.nidulans) between DNA repeats.

20 [0022] The invention further discloses host organisms transformed with the said vector.

[0023] The invention further discloses selection marker gene free recombinant microorganisms.

[0024] Specifically, the invention discloses organisms containing site-specifically introduced genes without any further foreign DNA being present. The method is therefore also suited for repeated modifications of the host genome, e.g. the sequential introduction of multiple gene copies at predetermined loci.

25 [0025] The invention provides a method for obtaining selection marker gene free recombinant strains comprising the following steps:

- a) integration into the genome of the strain of a desired DNA fragment and a selection marker,
b) selection of the recombinants,
30 c) deletion of the selection marker preferably using internal recombination between selection marker flanking repeats,
d) counter-selection based on the absence of the selection marker.

35 [0026] Although this is the preferred method for obtaining selection marker gene free recombinant strain, the invention also provides modifications of this method, for example:

The desired DNA fragment and the selection marker may be present on two different DNA molecules which are co-transformed. The selection marker does not necessarily integrate into the genome of the strain but may be present on an episomal DNA molecule which can be cured.

40 [0027] The present invention further illustrates that this marker gene can be deleted from the genome of the transformed organisms without leaving a trace i.e. DNA used for cloning.

[0028] The present invention discloses the use of the amdS gene from Aspergillus as a marker in bacteria and yeast.

45 [0029] The invention discloses also the use of the amdS gene for deleting a desired gene from the chromosome of a 'host' organism. Such modification techniques may be applied to filamentous fungi, yeasts and bacteria. In specific embodiments the following strains are employed Aspergilli, Trichoderma, Penicillium, Bacilli, E.coli, Kluyveromyces and Saccharomyces.

[0030] The method of the present invention provides recombinant strains with genomic modifications obtained by repeating the procedure with the same or other vectors.

50

55

Brief description of the drawingsAbbreviations used in the figures:5 Restriction enzymes and restriction sites:

[0031]

- 10 A = Apal; Ba = BamHI; B = BglII; Bs = BssHII; E = EcoRI;
H = HindIII; K = KpnI; N = NdeI; N = NotI; Ps = PstI;
P = PvuII; Sa = Sall; Sc = Scal; S = SmaI; Sn = SnaBI;
Spe = SpeI; Sp = SphI; Ss = SstII; Xb = XbaI; X = XhoI. Other:
T. = LAC4 terminator sequence
- 15 Figure 1: shows the restriction map of plasmid pamdS-1. This plasmid contains the cDNA of the amdS gene from A.nidulans.
- Figure 2: shows schematically the marker gene free deletion of the glaA locus from A.niger using the gene replacement vector pGBDEL4L. The essential part of the gene replacement vector pGBDEL4L contains the amdS gene under control of the gpdA promoter cloned between repeats (3'-non-coding region of the glaA gene).
- 20 Figures 3-9: show schematically the construction pathway of pGBDEL4L as further outlined in Example 1.
- Figure 10 A. KpnI digests of pGBDEL4L transformants #41 (lane 1), #24 (lane 2), #23 (lane 3) and #19 (lane 4) and the host strain A.niger CBS 513.88 (lane 5) and BamHI digests of pGBDEL4L transformants #41 (lane 6), #24 (lane 7), #23 (lane 8), #19 (lane 9) and the host strain A.niger CBS 513.88 (lane 10), probed with ³²P-labelled glaA promoter fragment and xylanase probe. B. KpnI digests of GBA-102 (lane 1) and the GBA-102 strains after fluoracetamide selection: GBA-107 (lane 2) and GBA-108 (lane 3) and BamHI digests of GBA-102 (lane 4) and the GBA-102 strains after fluoracetamide selection: GBA-107 (lane 5) and GBA-108 (lane 6), probed with ³²P-labelled glaA promoter fragment and xylanase probe.
- 25 Figure 11: A. Schematic presentation of BamHI and KpnI fragment lengths of the wild-type glaA locus in Aspergillus niger CBS 513.88.
B. Schematic presentation of BamHI and KpnI fragment lengths of the truncated glaA locus in transformant #19 (= GBA-102).
C. Schematic presentation of BamHI and KpnI fragment lengths of the truncated glaA locus in GBA-102 transformants after removal of the amdS gene (=GBA-107 and GBA-108).
- 30 Figure 12: A: shows schematically the integration of the glaA gene into the 3' non-coding region of truncated glaA locus of A.niger GBA-107.
B: shows the result of the internal recombination between the 3' glaA repeats, flanking the amdS gene.
- 40 Figures 13-24: show schematically the construction pathway of the integration vector pGBGLA30 as further outlined in Example 2.
- Figure 25: BglII digests of pGBGLA30 transformants #107-9 (lane 1), #107-7 (lane 2) and #107-5 (lane 3), the host strain A.niger GBA-107 (lane 4) and the parental strain A.niger CBS 513.88 (lane 5) and KpnI digests of pGBGLA30 transformants #107-9 (lane 6), #107-7 (lane 7) and #107-5 (lane 8), the host strain A.niger GBA-107 (lane 9) and the parental strain A.niger CBS 513.88 (lane 10), probed with ³²P-labelled glaA promoter fragment.
- 45 Figure 26: A: Schematic presentation of the KpnI and BglII fragment lengths of the wild-type glaA locus in Aspergillus niger CBS 513.88.
B: Schematic presentation of the KpnI and BglII fragment lengths of the truncated glaA locus in Aspergillus niger GBA-107.
C: Schematic presentation of the KpnI and BglII fragment lengths of the truncated glaA locus with a single copy pGBGLA30 integrated into the glaA 3'-non-coding region as in transformants #107-5 (= GBA-119) and #107-9 (= GBA-122).
D: Schematic presentation of the KpnI and BglII fragment lengths of the truncated glaA locus in GBA-119 and GBA-122 transformants after removal of the amdS gene (= GBA-120, GBA-121, GBA-123 and GBA-124).
- 50
- 55

- Figure 27: A: BglIII digests of A.niger CBS 513.88 (lane 10), GBA-107 (lane 9), GBA-119 (lane 8) and the GBA-119 strains after fluoracetamide selection: #AG5-7 (= GBA-120) (lane 5), #AG5-5 (= GBA-121) (lane 6) and #AG5-6 (lane 7); GBA-122 (lane 4) and the GBA-122 strains after fluoracetamide selection: #AG9-1 (= GBA-123) (lane 3), #AG9-2 (lane 2) and #AG9-4 (= GBA-124) (lane 1), probed with ³²P-labelled 3" glaA non-coding fragment.
- B: KpnI digests of A.niger CBS 513.88 (lane 10), GBA-107 (lane 9), GBA-119 (lane 8) and the GBA-119 strains after fluoracetamide selection: #AG5-7 (= GBA-120) (lane 5), #AG5-5 (= GBA-121) (lane 6) and #AG5-6 (lane 7); GBA-122 (lane 4) and the GBA-122 strains after fluoracetamide selection: #AG9-1 (= GBA-123) (lane 3), #AG9-2 (lane 2) and #AG9-4 (= GBA-124) (lane 1), probed with ³²P-labelled 3" glaA non-coding fragment.
- Figure 28: shows schematically the construction pathway of pGBGLA50.
- Figures 29-33: show schematically the construction pathway of pGBGLA53.
- Figure 34: shows schematically the construction of pGBamdS1.
- Figure 35: shows schematically the construction of pGBamdS2.
- Figure 36: shows schematically the construction of pGBamdS3.
- g shows schematically the construction of pGBamdS5.
- Figure 38: shows schematically the construction of pGBamdS6.
- Figure 39: shows schematically the construction of pPTLAC4 which was used in the construction of pGBamdS6.
- Figure 40: shows schematically the construction of pGBamdS7.
- Figure 41: HindIII digests of K.lactis CBS 683 (lane 1), K.lactis CBS 2360 (lane 2), the K.lactis CBS 683/pGBamdS1 transformants KAM-1 (lane 3), the K.lactis CBS 2360/pGBamdS1 transformant KAM-2 (lane 4) and the KAM-1 strains after fluoracetamide selection (lane 5,6) probed with a ³²P labelled LAC4 promoter fragment.
- Figure 42: A: BamHI digests of K.lactis CBS 683/pGBamdS3 transformants (lanes 1-3) probed with a ³²P-labelled LAC4 terminator fragment.
- B: BamHI digests of K.lactis CBS 683/pGBamdS5 transformants (lanes 1-5) and the host strain K.lactis CBS 683 (lane 6) probed with a ³²P-labelled LAC4 terminator fragment.
- Figure 43: BamHI digests of S.cerevisiae D273-10B (lane 1) and S.cerevisiae D273-10B/pGBamdS5 transformants (lanes 2-8) probed with a ³²P-labelled amdS fragment.
- Figure 44: HindIII digests of K.lactis CBS 2360 (lane 1), the K.lactis CBS 2360/pGBamdS6 transformant (lane 6) and strains from the K.lactis CBS 2360/pGBamdS6 transformant after fluoracetamide selection (lanes 2-5) probed with a ³²P-labelled LAC4 terminator fragment.
- Figure 45: restriction map of the Bacillus plasmid pBHA1.
- Figure 46: restriction map of the Bacillus plasmid pLNF.
- Figure 47: shows schematically the construction of pGBamdS21.
- Figure 48: shows schematically the construction of pGBamdS22.
- Figure 49: shows schematically the construction of pGBamdS23.
- Figure 50: shows schematically the construction of pGBamdS25.
- Figure 51: shows schematically the construction of pGBamdS41.

Detailed description of the invention

- [0032] The present invention discloses the use of a marker for selecting transformed host strains. The selection marker gene can be used on an episomal DNA vector. However, in the present invention, the marker gene is preferably integrated into the genome of the host strain. The advantage of the selection marker of the present invention is that it is a non-antibiotic dominant selection marker. Another advantage of the selection marker of the present invention is that it can be easily deleted from the transformed host organism. The deletion of the marker is based on dominant selection. As such the selection marker of the present invention is a dominant and bi-directional selection marker. To our knowledge it is the only selection-marker available which is bidirectional and dominant in both directions.
- [0033] In the present description we use the term 'selection marker gene'. With this term we mean the DNA coding for the marker protein in a functional form irrespective of whether it is the actual gene or the cDNA derived therefrom. The gene or cDNA is used dependent on the host organism and the expected splicing problems.
- [0034] In the present invention we use the term 'vector'. By this is intended any DNA molecule that can be introduced into a selected host irrespective of whether the vector integrates into the genome of the host cell or remains episomal. The vector contains a selectable marker gene functional in the selected host or can be co-transformed with another DNA molecule containing such a selection marker gene.

[0035] The present description uses the term 'desired heterologous or homologous genes or DNA fragments'. By this is intended a DNA fragment that may be obtained from the host strain or from another species or strain. The desired DNA fragment may contain any genetic element, parts thereof or combinations thereof, such as a gene (coding part or complete locus), a cDNA, a promoter, a terminator, an intron, a signal sequence, any regulatory DNA sequence or recognition sequence of DNA-binding proteins. The fragment may also be a DNA sequence that has been modified i.e. contains one or more nucleotide alterations (e.g. insertions, deletions, substitutions).

[0036] The present description further uses the term 'introduction' of a desired gene or DNA fragment. By this is intended an insertion, deletion, substitution of desired DNA sequences in a selected host cell.

[0037] The term 'genetic modification' used in the present invention refers to any modification of DNA sequences in a selected host cell which is the result of the introduction of any one of the above mentioned desired DNA fragments into the host cell, preferably by transformation or co-transformation.

[0038] In general all these genetic modifications can be performed using the method of the present invention with subsequent deletion of the selection marker gene. Due to the fact that the recombinant strain containing such a genetic modification does not contain the selection marker gene, the procedure of the present invention can be repeated, so that the modifications suggested above can be combined in the recombinant strain. Ultimately, the procedure of the present invention can be used repeatedly up to the point that a recombinant strain is obtained from which all undesired activities have been removed by deletion or inactivation of the corresponding genetic elements and which contains the desired activities at the desired levels by sequential introduction of the corresponding desired DNA fragments at desired copynumbers and preferably at desired and defined loci.

[0039] The A. nidulans acetamidase (amdS) gene allows A. nidulans to grow on acetamide as the sole N-source. For microorganisms that lack the possibility or only have a very limited capacity to use acetamide as the sole N-source the acetamidase gene can in principle be used as a selection marker provided that acetamide is taken up by the cells. The amdS gene has successfully been employed as a marker gene in Aspergilli (Kelly and Hynes (1985) EMBO J. 4, 475-479; Christensen et al. (1988) Bio/technology 6, 1419-1422), Penicillium (Beri and Turner (1987) Curr. Genet. 11, 639-641) and Trichoderma (Penttilä et al. (1987) Gene 61, 155-164).

[0040] The present invention for the first time discloses the use of the amdS gene from A. nidulans as a selection marker in organisms other than filamentous fungi. The use of this selection marker is disclosed in bacteria and yeasts. Specifically, the use is demonstrated in S. cerevisiae, in K. lactis in B. subtilis, in B. licheniformis and in E. coli. In view of the disclosed applicability of the selection marker in species selected from such diverse groups as fungi, yeasts and bacteria it is to be expected that the marker will also be applicable in other species pertaining to these groups. Use of this marker is therefore not restricted to the disclosed species.

[0041] The amdS gene from A. nidulans is capable of converting acetamide to ammonia and acetic acid. This property enables A. nidulans to grow on a medium containing acetamide as the sole N-source or C-source.

[0042] Another property of the amdS gene is that it is also able to convert fluoracetamide to ammonia and fluoracetic acid. Fluoracetic acid however is toxic to the cell. It is this property that forms the basis for another aspect of the present invention i.e. the production of marker gene free recombinant strains. The fluoracetamide converting property enables the counter-selection of transformed cells. The amdS gene is introduced into the host strain and integrated into the genome through homologous recombination. The transformed strains are selected on a medium containing acetamide as the sole N-source. Subsequently the selected strains are grown on a medium containing fluoracetamide and urea (or other preferably defined N-sources) as the sole N-sources. The surviving strains will have deleted the amdS gene.

[0043] The present invention uses the A. nidulans amdS gene as acetamidase marker gene. The relevant properties provided by the acetamidase encoded by the A. nidulans amdS gene, i.e. the ability to hydrolyse acetamide into ammonia and acetate as well as the ability to liberate fluoracetic acid from fluoracetamide, can also be provided by acetamidases from other sources. Use of an acetamidase marker gene is therefore not restricted to the A. nidulans amdS gene but includes any DNA sequence encoding a functional acetamidase.

[0044] The frequency of marker deletion is substantially increased by increasing the capacity of the gene for intra-chromosomal homologous recombination. To achieve this the amdS gene is preferably placed between DNA repeats. These repeats are not necessarily both present in the vector but may also be created by a single cross-over integration. Alternatively, one may omit flanking repeats and rely on other mechanisms for removal or inactivation of the marker gene. In that case, however, the outcome may be less predictable and may not result in removal but rather in mere inactivation of the marker gene.

[0045] The vector may be constructed in such a way that, after deletion of the marker gene, no extraneous foreign DNA (except the DNA of interest) remains in the chromosome of the host strain. The invention discloses a vector comprising:

- a) a desired DNA fragment destined for introduction into the host genome,
- b) optionally a DNA sequence that enables the vector to integrate (site-specifically) into the genome of the host strain,

c) a gene encoding an acetamidase (e.g. the amdS gene from A.nidulans) between DNA repeats.

[0046] Identical results may be obtained when the DNA-fragment destined for introduction into the host genome and the selectable marker gene (e.g. the acetamidase gene) are present on two different DNA molecules which are co-transformed, in which case the DNA molecule containing the selectable marker does not necessarily integrate into the host genome but may be present on an episomal DNA molecule which can be cured.

[0047] The sequences used for integration as mentioned under b) are used if site-specific (or better locus specific) integration is desired. If such a sequence is not present the vector nevertheless may integrate into the genome. This does not influence the ability to delete the selection marker gene.

[0048] The dominant counter-selection described above can be employed in the development of industrial production strains in various ways. The use of a dominant selection marker is especially advantageous in the development of improved production strains due to the fact that these strains are often diploid or polyploid.

[0049] The vector used for integration of the amdS gene preferably contains another gene of interest. The invention thus further enables the introduction of desired foreign or homologous genes or DNA elements in the host organisms of choice using the amdS gene as a marker. Subsequently the amdS gene is deleted. Preferably, the amdS and the desired genes or DNA elements are introduced site-specifically, whereafter the amdS gene is deleted.

[0050] Specifically, the invention discloses organisms containing site-specifically introduced genes without any further foreign DNA being present. The invention is used for integration of multiple copies of a desired gene or a DNA element at predetermined genomic loci.

[0051] The invention provides a method for obtaining selection of marker gene free recombinant strains comprising the following steps:

- integration of a desired gene or DNA element and a selection marker by homologous recombination between sequences incorporated in an expression cassette and sequences on the host chromosome,
- selection using the selection marker gene that is dominant,
- deletion of the selection marker gene using selection marker gene flanking regions,
- selection based on the absence of the selection marker gene (counter-selection).

[0052] The present invention further shows that this marker gene can be deleted from the chromosomes of the transformed organisms without leaving a trace i.e. DNA used for cloning. Moreover, the invention also shows that similar if not identical results can be obtained when the desired gene or DNA element and the selection marker are present on two different DNA molecules which are co-transformed.

[0053] Finally the invention discloses the use of the amdS gene for deleting a desired gene from the chromosome of a 'host' organism.

[0054] In view of the above, the method of the present invention is ideally suited for, but not limited to the cloning and expression of genes coding for proteins used in food, feed or pharmaceutical applications or genes involved in biosynthesis of antibiotics and other bio-active compounds, i.e. recombinant proteins and/or hosts-organisms that are subject to strict registration requirements.

[0055] Examples of such proteins are well known in the art and include chymosin, phytase, xylanases, amylases, cellulases and hemicellulases, cytokines and other pharmaceutical proteins, etc.

[0056] The same method is employed for deletion of genes coding for proteins that influence production levels of desired proteins again without leaving a marker gene in the genome. Such proteins include proteases which actively digest the desired products that are highly expressed in the host strain and that therefore have a reduced potential of producing and or secreting the desired proteins. A preferred method for the deletion of a given gene would use a DNA construct containing the following elements in a 5' to 3' order: sequences 5' of the gene to be deleted, directly fused to sequences 3' of the gene to be deleted, followed downstream by a functional selection marker gene (preferably an acetamidase gene), followed downstream by again sequences 3' of the gene to be deleted. In this case both sequences 3' of the gene to be deleted are chosen such that they form repeats flanking the selection marker gene. Transformation of this DNA construct and subsequent replacement of the chromosomal copy of the gene to be deleted by the DNA construct with cross-over points in the sequences 5' and 3' of the gene to be deleted results in deletion of the given gene. Subsequent intrachromosomal recombination between the repeats flanking the selection marker gene and counter-selection for these recombinants finally results in a selection marker free strain with the given gene deleted. The DNA construct used for this deletion can be constructed such that no foreign DNA or other traces of the genetic modification are left in the strain carrying the deletion.

[0057] The invention discloses selection marker gene free recombinant microorganisms. Such microorganisms can be organisms that, after the use of the disclosed technology, contain an extra copy of a desired gene (either homologous or heterologous). Such microorganisms can be re-transformed over and over by sequential application of the same technology to insert or delete additional copies of the same or other gene(s) of interest.

[0058] The microorganisms may also be characterized in that they have (a) predetermined gene(s) deleted or altered in any desired way.

[0059] The method of the present invention makes possible the fine-tuning of the production of desired proteins. This possibility is based on the ease with which repeated rounds of insertion and deletion can be performed. The method makes possible the insertion or deletion of a desired number of gene copies. Thus the proteins are produced in desired amounts and in desired ratios. This is especially useful for the production of mixtures of proteins or enzymes.

[0060] Whereas it is known that the acetamidase gene is capable of conversion of acetamide as the sole N-source in *Aspergillus* it is here shown that the acetamidase gene is easily deleted from the genome of transformed *Aspergilli*. To achieve this the amdS gene is cloned between direct repeats. In principle any direct repeat which allows for internal recombination can be employed. In the present examples this is demonstrated by cloning the amdS gene between 3' amyloglucosidase (glaA) non-coding DNA sequences.

[0061] It is shown that the amdS gene can be integrated and deleted upon plating on medium containing fluoracetamide and urea as N-sources.

[0062] It is further demonstrated that the amyloglucosidase gene can be deleted from the genome of *Aspergillus*. A replacement vector is constructed containing a part of the glaA promoter, a synthetic DNA sequence containing stop codons in all three reading frames, the amdS gene from *A. nidulans* under the control of the *A. nidulans* glyceraldehyde-3-phosphate dehydrogenase promoter and wherein the amdS gene is flanked by 3' glaA non-coding sequences. After transformation of *A. niger* the vector is integrated by double crossing-over thereby effectively replacing the amyloglucosidase gene. After selection for amdS activity the transformed strains are plated on fluoracetamide and urea. Selection resulted in strains wherein the amdS gene was deleted.

[0063] This example is an illustration of the possibility of using the amdS gene for deletion of a desired gene from the genome of an *Aspergillus* strain. Other genes can be eliminated or modified in a similar manner.

[0064] In a further example it is demonstrated that a gene can be inserted marker free at a predetermined site in the genome. An integration vector is constructed containing the *A. niger* glaA locus and the amdS gene flanked by two 3' glaA non-coding repeats.

[0065] The construct is shown to integrate at the amyloglucosidase locus. After selection on fluoracetamide the amdS gene is deleted. In this way a gene copy is integrated at a specific locus without leaving marker DNA.

[0066] It is evident from the above that the procedures described herein enable one of skill in the art to integrate or delete desired genes at predetermined loci without leaving selection marker DNA behind.

[0067] This method can be employed for gene amplification and gene replacement.

[0068] An especially important application would be the integration of desired genes. Followed by classical strain improvement whereafter the genes that may be adversely affected by the classical strain improvement techniques are replaced with fresh unaffected copies of the gene of interest without loss of expression level.

[0069] The system as described for *Aspergillus* above is expected to give the same results when other fungal strains are employed, which are known to be incapable of growth on acetamide as the sole N-source. The use of the amdS gene as a selection marker has been described for among others *Penicillium* and *Trichoderma*. Moreover, the amdS gene can even be used in filamentous fungi which are capable of using acetamide as sole N-source albeit poorly. In this case the background of poorly growing untransformed cells can be repressed by the inclusion of CsCl in the selection media (Tilburn, J. et al. (1983) *Gene*, 26, 205-221). Hence the system is expected to be applicable to filamentous fungi in general.

[0070] In one embodiment of the present application it is surprisingly demonstrated that the *A. nidulans* amdS gene can be used as a selection marker in *K. lactis*. In this Example it is shown that two different *K. lactis* strains cannot grow on acetamide as the sole N-source. The two *K. lactis* strains are plated on YCB medium which is

- a) complete but without N-source,
- b) as a) but with acetamide,
- c) as a) but with ammonium sulphate.

[0071] It is shown that the strains do not grow on the medium under b) but do grow on medium under c). Hence provided that the acetamide is taken up by the yeast cells and that the amdS gene can be expressed in *K. lactis* the system is applicable in yeasts also at least as a selection marker. Concerning the counter-selection using fluoracetamide some further requirements have to be met. Fluoracetate is toxic when activated by the enzyme acetyl-CoA-synthetase. Prerequisites for the fluoracetamide counter-selection to also work on amdS⁺ yeasts are therefore

- 1) fluoracetamide should not be toxic for amdS⁺ yeasts,
- 2) the yeast cell wall and plasmamembrane should be permeable to fluoracetamide and
- 3) the enzyme acetyl-CoA-synthetase should be active.

[0072] To test this the amdS gene was cloned in K. lactis.

[0073] To avoid any potential splicing problems of the A.nidulans amdS gene in K.lactis the amdS cDNA from A. nidulans was cloned as shown in the Experimental section.

[0074] Subsequently the amdS was cloned downstream of a yeast promoter (LAC4, ADH1, K1EF) in a vector containing another marker (phosphotransferase-G418). This cloning is described in Example 8. The vectors containing both the G418 marker and the amdS gene were selected using the G418 marker and were then used to optimize selection conditions for the amdS⁺ phenotype.

[0075] Direct selection of K.lactis is shown in another embodiment of the present invention and for S.cerevisiae direct selection is shown in Example 11.

[0076] Subsequently it is demonstrated that counter-selection can be employed on the transformed yeast strains to remove the amdS gene.

[0077] The amdS gene system is used for both marker gene free insertion and marker gene free deletion of a gene in yeast.

[0078] In a further embodiment the lactase gene is deleted from K.lactis whereas in Example 14 a copy of the chymosin gene is inserted into the K.lactis genome.

[0079] The genes used here for insertion and deletion are only used as examples. The same technology can be applied using other genes or DNA elements. As mentioned before the DNA fragments used for insertion or deletion can be mutated genes, promoter sequences, regulatory sequences etc. In all cases it is possible to insert or delete these sequences at desired genomic sites and in desired numbers, without leaving a marker gene behind.

[0080] The feasibility of the use of this system in other yeast strains is evident.

[0081] As a first step for use of the system of the present invention in bacteria it is shown in Example 15 that Bacillus subtilis and E.coli cannot grow on acetamide as the sole N-source.

[0082] Example 16 describes the vectors that have been constructed for use in Bacillus and E.coli.

[0083] It is demonstrated in Examples 17 and 18 that the amdS gene can be effectively used in Bacillus and E.coli as selection marker, whereas Example 19 demonstrate the fluoracetamide counter-selection of bacterial amdS⁺ transformants.

[0084] The advantages of the system of the present invention are manifold. The most striking advantages are given below:

- It is demonstrated that the amdS system is universally applicable (plant cells, animal cell, yeasts, bacteria and filamentous fungi etc.), requiring only that the host in question cannot or only poorly grow on acetamide as sole C- or N-source but can utilize either acetate or ammonia as sole C- or N-source, respectively.
- The amdS system represents the only bi-directional and dominant selection system. This feature is extremely convenient for use in poly- or aneuploid strains which often is the case with natural isolates and/or industrial strains.
- After classical strain improvement any mutated copies of the desired gene can be easily replaced by unmutated copies by gene replacement due to the fact that the desired genes have been integrated at well-defined loci. The genes are thus replaced with unmutated genes without affecting the expression level.
- Due to the ability to introduce multiple integrations at well-defined and therefore non-random loci one can be assured that no undesirable traits arise in the strain upon gene amplification.
- The growing concern about the release of various selection markers in the environment is overcome by the presented system. No selection marker gene or other unnecessary or undesired DNA sequences need to present in the production strains after introduction of the desired genes or other genetic modifications.

Experimental

General molecular cloning techniques

[0085] In the examples described herein, standard molecular cloning techniques such as isolation and purification of nucleic acids, electrophoresis of nucleic acids, enzymatic modification, cleavage and/or amplification of nucleic acids, transformation of E.coli, etc., were performed as described in the literature (Sambrook et al. (1989) "Molecular Cloning: a laboratory manual", Cold Spring Harbour Laboratories, Cold Spring Harbour, New York; Innis et al. (eds.) (1990) "PCR protocols, a guide to methods and applications" Academic Press, San Diego). Synthesis of oligo-deoxynucleotides and DNA sequence analysis were performed on an Applied Biosystems 380B DNA synthesizer and 373A DNA sequencer, respectively, according to the user manuals supplied by the manufacturer.

Transformation of A.niger

[0086] Transformation of A.niger was performed according to the method described by Tilburn, J. et.al. (1983) Gene

26, 205-221 and Kelly, J. & Hynes, M. (1985) EMBO J., 4, 475-479 with the following modifications:

- spores were grown for 16 hours at 30°C in a rotary shaker at 300 rpm in Aspergillus minimal medium. Aspergillus minimal medium consists of the following components: Per liter: 6 g NaNO₃; 0.52 g KCl; 1.52 g KH₂PO₄; 1.12 ml 4M KOH; 0.52 g MgSO₄·7H₂O; 10 g glucose; 1 g casamino acids; 22 mg ZnSO₄·7H₂O; 11 mg H₃BO₃; 5 mg FeSO₄·7H₂O; 1.7 mg CoCl₂·6H₂O; 1.6 mg CuSO₄·5H₂O; 5 mg MnCl₂·4H₂O; 1.5 mg Na₂MoO₄·2H₂O; 50 mg EDTA; 2 mg riboflavin; 2 mg thiamine.HCl; 2 mg nicotinamide; 1 mg pyridoxine.HCl; 0.2 mg panthotenic acid; 4 µg biotin; 10 ml Penicillin (5000 IU/ml)/Streptomycin (5000 UG/ml) solution (Gibco).
- only Novozym 234 (Novo Industri), and no helicase, was used for formation of protoplasts;
- after protoplast formation (60-90 minutes), KC buffer (0.8 M KCl, 9.5 mM citric acid, pH6.2) was added to a volume of 45 ml. and the protoplast suspension was centrifuged at 2500 g at 4°C for 10 minutes in a swinging-bucket rotor. The protoplasts were resuspended in 20 ml. KC buffer. Then, 25 ml of STC buffer (1.2 M sorbitol, 10 mM Tris-HCl pH7.5, 50 mM CaCl₂) was added and subsequently the protoplast suspension was centrifuged at 2500 g at 4°C for 10 minutes in a swinging-bucket rotor, washed in STC-buffer and resuspended in STC-buffer at a concentration of 10⁸ protoplasts/ml;
- to 200 µl of the protoplast suspension the DNA fragment, in a volume of 10 µl in TE buffer (10 mM Tris-HCl pH7.5, 0.1 mM EDTA), was added and subsequently 100 µl of a PEG solution (20% PEG 4000 (Merck), 0.8 M sorbitol, 10 mM Tris-HCl pH7.5, 50 mM CaCl₂);
- after incubation of the DNA-protoplast suspension at room temperature for 10 minutes, 1.5 ml PEG solution (60% PEG 4000 (Merck), 10 mM Tris-HCl pH7.5, 50 mM CaCl₂) was added slowly, with repeated mixing of the tubes. After incubation at room temperature for 20 minutes, the suspensions were diluted with 5 ml STC buffer, mixed by inversion and centrifuged at 2000 g at room temperature for 10 minutes. The protoplasts were resuspended gently in 1 ml 1.2 M sorbitol and plated onto selective regeneration medium consisting of Aspergillus minimal medium without riboflavin, thiamine.HCl, nicotinamide, pyridoxine.HCl, panthotenic acid, biotin, casamino acids and glucose but with 10 mM acetamide as the sole nitrogen source, 1 M sucrose, solidified with 2% bacteriological agar #1 (Oxoid, England). Following growth for 6-10 days at 30°C, the plates were replica plated onto selective acetamide plates consisting of Aspergillus selective regeneration medium with 2% glucose instead of sucrose and 1.5% agarose instead of agar. Single transformants were isolated after 5-10 days of growth at 30°C.

30 Transformation of A. oryzae

[0087] Transformation of A. oryzae was performed according to the method described by Christensen, T. et al. in European Patent Application 0 238 023 A2.

35 Transformation of T. reesei

[0088] Transformation of T. reesei was performed according to the method described by Penttillä M., Knowles, J. (1987) Gene 61 155-164.

40 Transformation of P. chrysogenum

[0089] The Ca-PEG mediated protoplast transformation procedure is used. Preparation of protoplasts and transformation of P.chrysogenum was performed according to the method described by Gouka et al., Journal of Biotechnology 20(1991), 189-200 with the following modifications:

- After transformation, the protoplasts were plated onto selective regeneration medium plates consisting of Aspergillus minimal medium, osmotically stabilized with 1.2 M sucrose, containing 0.1% acetamide as sole nitrogen source and solidified with 1.5% bacteriological agar #1 (Oxoid, England).
- After 5-8 days of incubation at 25°C transformants appeared.

50 Transformation of K.lactis

[0090] The yeast K.lactis was transformed using the lithium acetate procedure described by Ito H. et al. (1983) J. Bacteriol. 153, 163-168 with the following modifications:

- For transformation a K.lactis culture was taken with an OD₆₁₀ between 0.5 and 1.0.
- After the 5 minutes heatshock of the transformed cell suspensions, 1 ml YEPD/YNB (1% yeast-extract, 2% Bacto-peptone, 2% glucose and 0.17% Yeast Nitrogen Base w/o amino acids (YNB; Difco) was added and the cell-

suspensions were incubated at 30°C in a shaker incubator for 150 - 180 minutes.

- After the above mentioned incubation (at 30°C for 150 - 180 minutes), the cell-suspensions were centrifuged at 2000 g at room temperature for 5 minutes and plated on YEPD/G418 double layer medium solidified with 2% Bactoagar (Difco). YEPD/G418 double layer plates were prepared as followed: 10 minutes prior to plating of the cell-suspensions 15 ml YEPD agar (1% yeast-extract, 2% Bacto-peptone, 2% glucose solidified with 2% Bactoagar (Difco)) without G418 was poured onto 15 ml YEPD agar, which contained 50 µg G418/ml. This results in YEPD/G418 double layer plates which contain 25 µg G418/ml after diffusion of the antibiotic. The YEPD/G418 double layer plates contained 25 µg G418/ml or 100 µg G418/ml in case of strains *K.lactis* CBS 683 or CBS 2360, respectively.

Isolation of DNA from *Aspergillus*, *Trichoderma*, *Penicillium* and yeast

[0091] The isolation of DNA from *Aspergillus* and *Trichoderma* was performed according to the procedure as described by Yelton, et al. (1984), Proc. Natl. Acad. Sci. **81**, 1470-1474.

[0092] The isolation of DNA from *Penicillium* was performed according to the procedure described by Kolar et al., Gene **62** (1988), 127-134.

[0093] The isolation of DNA from *K.lactis* or *S.cerevisiae* was performed according to the procedures described by Fujimura and Sakuma (1993), Biotechniques **14**, 538.

Bacillus transformation and DNA-isolation

[0094] Transformation of the different *Bacillus* species as well as isolation of plasmid or chromosomal DNA from these species was performed as described by Bron (1990) "Plasmids" In: Molecular Biological Methods for *Bacillus*, Harwood, CR and Cutting, SM, eds., series Modern Microbiological Methods, John Wiley & Sons, Chichester, UK.

[0095] For the transformation of *B.subtilis* BS-154 (CBS 363.94) competent cells were used and for the transformation of *B.licheniformis* T5 (CBS 470.83) protoplast transformation was used. In the case of neomycin selection a concentration of 20 µg/ml was used. For acetamide selection of *B.subtilis* transformants, minimal medium agar was used in which casamino acids and yeast extract were replaced by 20 mM acetamide. For acetamide selection of *B.licheniformis* transformants, protoplast regeneration medium was used in which ammonium sulphate was replaced by 20 mM acetamide.

Removal of the *amdS* selection marker

[0096] The *amdS* marker in most examples relating to *Aspergillus*, *Trichoderma* and *Penicillium* is cloned between repeats consisting of a part of the 3' non-coding region of amyloglucosidase gene. Removal of the *amdS* selection marker is achieved either by internal recombination between the 3' *glaA* non-coding repeats that flank the *amdS* selection marker or by homologous recombination between the repeats that are created by integration via a single cross-over event. Selection of cells that have lost the *amdS* selection marker is achieved by growth on plates containing fluoracetamide. Cells harbouring the *amdS* gene metabolize fluoracetamide to ammonium and fluoracetate which is toxic to the cell. Consequently, only cells that have lost the *amdS* gene are able to grow on plates containing fluoracetamide.

[0097] In case of removal of the *amdS* marker from *Aspergillus* transformants, spores from these transformants were plated onto selective regeneration medium (described above) containing 32 mM fluoracetamide and 5 mM ureum instead of 10 mM acetamide, 1.1% glucose instead of 1M sucrose and 1.1% instead of 2% bacteriological agar #1 (Oxoid, England). After 7-10 days of growth at 35°C single colonies were harvested and plated onto 0.4% potato dextrose agar (Oxoid, England). In case of removal of the *amdS* marker from *Trichoderma* transformants, spores of these transformants were plated onto non selective minimal medium plates (per liter: 20 g. glucose, 5 g. (NH₄)₂SO₄, 15 g. KH₂PO₄, 0.6 g. MgSO₄, 0.6 g. CaCl₂, 0.005 g. FeSO₄·7H₂O, 0.0016 g. MnSO₄·H₂O, 0.0014 g. ZnSO₄·7H₂O, 0.002 g. CoCl₂; pH5.5) supplemented with 10 mM fluoracetamide. After 5-10 days at 30°C, colonies were harvested and plated onto 0.4% potato dextrose agar (Oxoid, England).

[0098] In case of removal of the *amdS* marker from *Penicillium* transformants, spores from these transformants were plated on selective medium plates consisting of *Aspergillus* minimal medium with 10 mM fluor-acetamide and 5% glucose, solidified with 1.5% bacteriological agar #1 (Oxoid, England). After 5-10 days of growth at 25°C resistant colonies appeared.

Determination of glucoamylase production by *A.niger* transformants

[0099] Of recombinant and control *A.niger* strains spores were collected by plating spores or mycelia onto PDA-

plates (Potato Dextrose Agar, Oxoid), prepared according to the supplier's instructions. After growth for 3-7 days at 30°C spores were collected after adding 0,01% Triton X-100 to the plates. After washing with sterile water approximately 10⁷ spores of selected transformants and control strains were inoculated into shake flasks, containing 20 ml of liquid preculture medium containing per litre: 30 g maltose.H₂O; 5 g yeast extract; 10 g hydrolysed casein; 1 g KH₂PO₄; 0.5 g MgSO₄.7H₂O; 3 g Tween 80; 10 ml Penicillin (5000 IU/ml)/Streptomycin (5000 UG/ml); pH 5.5. These cultures were grown at 34°C for 20-24 hours. 5-10 ml of this culture was inoculated into 100 ml of fermentation medium containing per litre: 70 g maltodextrines; 25 g hydrolysed casein; 12.5 g yeast extract; 1 g KH₂PO₄; 2 g K₂SO₄; 0.5 g MgSO₄.7H₂O; 0.03 g ZnCl₂; 0.02 g CaCl₂; 0.01 g MnSO₄.4H₂O; 0.3 g FeSO₄.7H₂O; 10 ml penicillin (5000 IU/ml)/Streptomycin (5000 UG/ml); adjusted to pH 5.6 with 4 N H₂SO₄. These cultures were grown at 34°C for 5-10 days. Samples were taken for the analysis of the glucoamylase production at different time points during fermentation. Fermentation broth samples were centrifuged (10 minutes, 10.000xg) and supernatants collected.

[0100] The glucoamylase activity was determined by incubating 10 µl of a six times diluted sample of the culture supernatant in 0.032 M NaAc/HAc pH4.05 with 115 µl of 0.2% (w/v) p-Nitrophenyl α-D-glucopyranoside (Sigma) in 0.032 M NaAc/HAc pH 4.05. After a 30 min incubation at room temperature, 50 µl of 0.3 M Na₂CO₃ was added and the absorption at a wavelength of 405 nm was measured. The A_{405nm} is a measure for the AG production.

Cloning of the amdS cDNA

[0101] The *A.nidulans* *amdS* gene contains three small introns (Corrick et al. (1987) Gene 53, 63-71). In order to avoid problems caused by incorrect splicing of these introns in yeast or lack of splicing in bacteria, we have used an *amdS* cDNA for expression in yeasts and bacteria. Cloning of the *amdS* cDNA from an *A.nidulans* polyA⁺ RNA preparation has been described by Corrick et al. ((1987), Gene 53, 63-71). In this example we have used the *A.niger* NRRL 3135 transformant #4, which is transformed by multiple copies of the *A.nidulans* *amdS* gene containing plasmid pAF2-2S (van Hartingsveld et al (1993). Gene 127, 87-94). Total RNA was isolated by a direct LiCl precipitation according to a procedure modified from Auffray et al. ((1980) Eur.J.Biochem. 107, 303-314). *A.niger* spores were allowed to germinate and were grown overnight at 37°C in a minimal medium (Cove (1966) Biochim. Biophys. Acta 113, 51-56) supplemented with glucose as carbon source and with acetamide as sole nitrogen source. Mycelium was obtained and dried by filtration and subsequently frozen with liquid nitrogen to be grounded. The powder was dispersed in 3M LiCl, 6M urea at 0°C and maintained overnight at 4°C. Total cellular RNA was obtained after centrifugation at 16.000 g for 30 minutes and two successive extractions with phenol/chloroform/isoamylalcohol (50:48:2). The RNA was precipitated with ethanol and dissolved in 1 ml 10 mM Tris-HCl (pH7.4), 0.5% SDS. For polyA⁺ selection the total RNA sample was heated for 5 minutes at 65°C and subsequently applied to an oligo(dT)-cellulose column. After several washes with a solution containing 10 mM Tris-HCl pH 7.4, 0.5% SDS and 0.5 M NaCl, the polyA⁺ RNA was collected by elution with 10 mM Tris-HCl pH 7.4 and 0.5% SDS and precipitated with ethanol. Approximately 5 µg of the polyA⁺ mRNA was used as template for reverse transcription primed with oligo(dT) primers. The reaction mixture (50 mM Tris-HCl pH 7.6, 10 mM DTT, 6 mM MgCl₂, 80 mM KCl, 0.2 mM each dNTP and 0.1 mg BSA/ml) was incubated for 30 minutes at 37°C with 500 units Murine MLV reverse transcriptase (BRL) and 75 units RNase inhibitor (Promega) in a volume of 100 µl. Another 200 units of reverse transcriptase were added and the reaction was continued for 30 minutes. The mixture was extracted with chloroform and precipitated with ethanol in the presence of 0.25 M ammonium acetate. This mixture of first strand cDNAs was used as template in a subsequent Polymerase Chain Reaction (PCR) to amplify the *amdS* cDNA. The genomic *amdS* sequence was used to design 2 synthetic oligonucleotides that were used as primers in this PCR:

AB3100 (SEQ ID NO: 1):

5'-CTAATCTAGAATGCCTCAATCCTGAA-3' (an *amdS*-specific sequence from nucleotide -3 to +16 preceded by an XbaI site and 4 additional nucleotides).

AB3101 (SEQ ID NO: 2):

5'-GACAGTCGACAGCTATGGAGTCACCACA-3' (an *amdS*-specific sequence positioned downstream of the *amdS* stopcodon from nucleotides 1911 to 1884 flanked by an additional SalI site).

[0102] The PCR reaction was performed using 10 % of the cDNA mixture as template and 0.1 µg of each of the oligos AB3100 (SEQ ID NO: 1) and AB3101 (SEQ ID NO: 2) as primer. After denaturation (7 minutes at 100 °C) and addition of 1.3 units Taq-polymerase the reaction mixture was subjected to 25 amplification cycles (each cycle: 2 minutes at 94°C, 2 minutes at 55°C and 3 minutes at 72°C). In the last cycle the extension step was longer (7 min.) to allow synthesis of full-length fragments. The obtained DNA fragment was digested with XbaI and SalI and subcloned into the XbaI/SalI sites of pUC18. The resulting plasmid was designated pamdS-1 (see Figure 1). Restriction analysis of the plasmid pamdS-1 confirmed the absence of introns and the correct fusion of exons in the *amdS* cDNA.

Example 1

Marker gene free deletion of an *A.niger* gene by using the *amdS* gene

[0103] In this example a genomic target gene in *A.niger* will be replaced by transforming *A.niger* with a replacement vector which integrates into the *A.niger* genome via a double cross-over homologous recombination. The replacement vector comprises a DNA region homologous to the target locus interrupted by a selectable marker gene flanked by DNA repeats.

[0104] In this example plasmid pGBDEL4L is used to delete the *glaA* coding region and a (proximal) part of the *glaA* promoter region. This vector comprises a part of the *A.niger glaA* genomic locus, wherein the *glaA* coding sequences as well as a part of the *glaA* promoter sequences are replaced by the *A.nidulans amdS* gene under the control of *A.nidulans gpdA* promoter as selection marker flanked by 3'-untranslated *glaA* sequences as direct repeats. Transformation of *A.niger* with this vector directs replacement of the *glaA* gene by the *amdS* marker gene. By performing the fluoracetamide counter-selection on these transformants as described in the experimental procedures, the *amdS* marker gene will be deleted properly by an internal recombination event between the 3'*glaA* DNA repeats, resulting in a marker gene free Δ *glaA* recombinant strain, possessing finally no foreign DNA sequences at all (for a schematic view, see Figure 2).

Short description of the *glaA* gene replacement vector pGBDEL4L

[0105] The gene replacement vector pGBDEL4L contains 5'-part of the *A.niger* amyloglucosidase (*glaA*) promoter region, a synthetic DNA sequence of 16 bp providing stopcodons in all three reading frames, the *A.nidulans* acetamidase (*amdS*) gene under control of the *A.nidulans* glyceraldehyde-3-phosphate dehydrogenase (*gpdA*) promoter, flanked at both sides by 3' *glaA* non-coding sequences.

Construction pathway of pGBDEL4L

[0106] In order to obtain the final deletion vector pGBDEL4L several subclones of the *glaA* locus were derived first. A schematic view is presented in figure 3. The *glaA* locus of *A.niger* was molecularly cloned and described previously (EP 0 463 706 A1). Plasmid pAB6-1 contains the entire *glaA* locus from *A.niger* on a 15.5 kb *HindIII* fragment cloned in the *HindIII* site of pUC19 (Yanisch-Perron *et al.*, Gene 33 (1985) 103-119, and is obtainable from e.g. Boehringer Mannheim, Germany). pAB6-1 was digested with *EcoRI* and the 1.8 kb *EcoRI* DNA fragment just upstream of the *glaA* gene was isolated by agarose gel electrophoresis and ligated into pUC19 digested with *EcoRI* and subsequently transferred to *E.coli* and molecularly cloned. The resulting plasmid was designated pAB6-3 (Figure 3A). To construct plasmid pAB6-4, which is another subclone of pAB6-1, pAB6-1 was digested with *HindIII* and *BglII*. The 4.6 kb sized DNA fragment comprising the *glaA* promoter and a part of the *glaA* coding sequence was isolated by agarose gel electrophoresis and ligated into pUC19 which was digested prior with *HindIII* and *BamHI* (Figure 3B). As a result the *BamHI* as well as the *BglII* sites in pAB6-4 were destroyed appropriately by this cloning procedure.

[0107] Subsequently, after digesting plasmid pAB6-4 with *HindIII* and *EcoRI* and filling in the 5' sticky ends using *E.coli* DNA polymerase, the 1.8 kb *glaA* promoter DNA fragment was isolated by agarose gel electrophoresis, ligated into pAB6-3 which was partially digested with *EcoRI* and treated with *E.coli* DNA polymerase to generate blunt ends, the ligation mixture was transferred to *E.coli* for molecular cloning. The derived plasmid (designated pAB6-31) contains a 3.6 kb *glaA* promoter fragment with a destroyed *EcoRI* site in the middle, but still possessing the *EcoRI* site (now unique in this DNA fragment) just upstream of the *glaA* ATG initiation site (Figure 4).

[0108] The *A.nidulans amdS* gene used herein is located on an approximately 4 kb sized *EcoRI-KpnI* fragment in plasmid pGW325 (Wernars *et al.*, thesis (1986) Agricultural University, Wageningen, The Netherlands). This *EcoRI-KpnI* DNA fragment containing the *amdS* gene, flanked by its own regulatory sequences, was molecularly cloned into the appropriate sites of pUC19 as described by Verdoes *et al.* (Transgenic Res. 2 pp 84-92, 1993) resulting in pAN4-1. pAN4-1 was digested with *EcoRI* and *KpnI*, the 4 kb sized DNA fragment containing the *amdS* gene was isolated by agarose gel electrophoresis, ligated into pAB6-31 digested with *EcoRI* and *KpnI* and the ligation mixture was transferred to *E.coli* for molecular cloning. The obtained plasmid was designated pAB6S (Figure 5) and contains a 3.8 kb *glaA* promoter DNA fragment and the 4 kb *amdS* fragment.

[0109] Plasmid pAB6S was first partially digested with *Sall*, and ligated to the synthetic derived oligonucleotide TN0001 (SEQ ID NO: 3) having the following sequence:

TN0001 (SEQ ID NO: 3): 5' TCGATTAAGTAGTTAA 3'

and secondly digested with EcoRI. The DNA fragment comprising the pUC19, the glaA promoter and the amdS gene sequences was purified and isolated by agarose gel electrophoresis. From plasmid pAB6-1, digested with Sall, the 2.2 kb 3' flanking glaA DNA fragment was isolated as well by agarose gel electrophoresis and ligated to the above mentioned synthetic oligonucleotide, treated with T4 polynucleotide kinase, subsequently digested with EcoRI and ligated to the above mentioned DNA fragment isolated of pAB6S. The DNA ligation mixture was transferred to E. coli and molecular cloned. The derived plasmid was designated pGBDEL1 and is shown in Figure 6. By this procedure simultaneously the Sall restriction site was destroyed and stopcodons in all reading frames were introduced.

[0110] To obtain an approximately 1 kb large DNA fragment, containing 3' glaA non-coding DNA sequences positioned just downstream the stop codon of the glaA gene and flanked by suitable restriction sites, a PCR amplification was performed. In this PCR amplification, the plasmid pAB6-1 was used as template and as primers two synthetical derived oligonucleotides:

Oligo AB2154 (SEQ ID NO: 4):

5' AACCATAGGGTCGACTAGACAATCAATCCATTTTCG 3'

(a 3' glaA non-coding sequence just downstream of the stopcodon) and

Oligo AB2155 (SEQ ID NO: 5):

5' GCTATTTCGAAAGCTTATTCATCCGGAGATCCTGAT 3'

(a 3' glaA non-coding sequence around the EcoRI site approx. 1 kb downstream of the stopcodon).

[0111] The PCR was performed as described by Saiki et al. (Science 239, 487-491, 1988) and according to the supplier of TAQ-polymerase (Cetus). Twenty five amplification cycles (each 2 minutes at 55 °C; 3 minutes at 72 °C and 2 minutes at 94 °C) were performed in a DNA-amplifier (Perkin-Elmer/Cetus). The 1 kb amplified DNA fragment was digested with HindIII and Sall, purified by agarose gel electrophoresis, ethanol precipitated and subsequently cloned into the HindIII and Sall restriction sites of pGBDEL1. The thus obtained plasmid was designated pGBDEL2 (Figure 7A,B).

[0112] To obtain the final glaA gene replacement vector pGBDEL4L, the amdS promoter region in pGBDEL2 was exchanged by the stronger A.nidulans gpdA promoter. Fusion of the gpdA promoter sequence to the coding sequence of the amdS gene was performed by the Polymerase Chain Reaction (PCR) method. For this PCR fusion two different templates were used: plasmid pAN7-1 (Punt et al., Gene 56, 117-124, 1987) containing the E.coli hph gene under control of the A.nidulans gpdA promoter and the A.nidulans trpC terminator and plasmid pAN4-1, containing the A.nidulans amdS gene under control of its own regulatory sequences. As primers four synthetic oligonucleotides were used, possessing the following sequences:

Oligo AB 2977 (SEQ ID NO: 6):

5' TATCAGGAATTCGAGCTCTGTACAGTGACC 3'

(a 5' gpdA promoter specific oligo nucleotide, positioned at approximately 880 bp upstream of the ATG startcodon of the E. coli hph gene)

Oligo AB2992 (SEQ ID NO: 7):

5' GCTTGAGCAGACATCACCATGCCTCAATCCTGGGAA 3'

Oligo AB2993 (SEQ ID NO: 8):

5' TTCCAGGATTGAGGCATGGTGATGTCTGCTCAAGC 3'

(both sequences are complementary to each other and contain 18 bp of the 3' end of the gpdA promoter and 18 bp of the 5' part of the amdS coding region)

Oligo AB2994 (SEQ ID NO: 9):

5' CTGATAGAATTTCAGATCTGCAGCGGAGGCCTCTGTG 3'

(an amdS specific sequence around the BglII site approximately 175 bp downstream of the ATG initiation codon)

[0113] To fuse the 880 bp gpdA promoter region to the amdS coding sequence two separate PCR's were carried out: the first amplification with pAN7-1 as template and the oligo nucleotides AB 2977 (SEQ ID NO: 6) and AB2993 (SEQ ID NO: 8) as primers to amplify the 880 bp DNA fragment comprising the gpdA promoter flanked at the 3' border by 18 nucleotides complementary to the 5' end of the amdS gene, and the second PCR reaction with pAN4-1 as template and the oligo nucleotides AB2992 (SEQ ID NO: 7) and AB2994 (SEQ ID NO: 9) as primers to amplify a 200 bp sized DNA fragment comprising the 5' part of the amdS gene flanked at the 5' border by 18 nucleotides complementary to the 3' end of the gpdA promoter. A schematic view of these amplifications is presented in Figure 8A. The two fragments generated were subsequently purified by agarose gel electrophoresis, ethanol precipitated and used as templates in a third PCR reaction with oligo nucleotides AB 2977 (SEQ ID NO: 6) and AB2994 (SEQ ID NO: 9) as primers. The resulting DNA fragment was digested with EcoRI, purified by agarose gel electrophoresis and ethanol precipitation, and cloned into the EcoRI site of pTZ18R (United States Biochemicals). The resulting plasmid was designated pGBGLA24 (Figure 8B).

[0114] To exchange the amdS promoter sequence in pGBDEL2 by the gpdA promoter sequence, the approximately 1 kb sized EcoRI/BglII DNA fragment of pGBGLA24 was isolated by agarose gel electrophoresis after digestion with the appropriate restriction enzymes and ligated into the EcoRI and BglII sites of pGBDEL2. The resulting glaA gene replacement vector was designated pGBDEL4L (Figure 9).

Deletion of glaA promoter and coding sequences in A.niger

[0115] Prior to transformation of A.niger with pGBDEL4L, the E.coli sequences were removed by HindIII and XhoI digestion and agarose gel electrophoresis. The A.niger strain CBS 513.88 (deposited October 10, 1988) was transformed with either 2.5, 5 or 10 µg DNA fragment by procedures as described in experimental procedures using acetamide as sole N-source in selective plates. Single A.niger transformants were purified several times onto selective acetamide containing minimal plates. Spores of individual transformants were collected by growing for about 5 days at 30°C on 0.4% potato-dextrose (Oxoid, England) agar plates. Southern analyses were performed to verify the presence of the truncated glaA locus. High molecular weight DNA of several transformants was isolated, digested with BamHI and KpnI and subsequently fractionated by electrophoresis on a 0.7% agarose gel. After transfer to nitrocellulose filters, hybridization was performed according to standard procedures using two ³²P-labelled probes: a XhoI/SalI glaA promoter fragment isolated from plasmid pAB6-4 (described above, Figure 3A) and a probe recognizing endogenous xylanase sequences (European Patent Application. 0 463 706 A). The results of only 4 transformants (#19, #23, #24, #41) and the control strain A.niger CBS 531.88 are shown as examples in Figure 10A. For a better understanding of this autoradiograph, a schematic presentation is presented in Figure 11 showing the size of the hybridizing fragments in intact and truncated glaA loci.

[0116] Characteristic for the intact glaA locus is a 3.5 kb hybridizing fragment in a BamHI digest and a 4.5 kb hybridizing fragment in a KpnI digest (see figure 11A). In a truncated glaA locus, the 3.5 kb BamHI hybridizing fragment and the 4.5 kb KpnI hybridizing fragment are absent and replaced by a 5.5 kb BamHI hybridizing fragment and a 6.3 kb KpnI hybridizing fragment. In this example, as can be seen in Figure 10A, transformant #19 shows the expected pattern of a truncated glaA locus (Figure 11B). This transformant was designated GBA-102.

[0117] No replacement of the glaA gene had occurred in the other transformants. The poorly hybridizing bands: 4, 8 and 15 kb in the KpnI digest and 7 and 12 kb in the BamHI digest, refer to the xylanase sequences as internal control.

Removal of the amdS gene from A.niger GBA-102 by counter-selection on fluoracetamide containing plates.

[0118] The amdS gene in the transformant A.niger GBA-102 was removed again as described in the Experimental section. The removal of the amdS selection marker gene in only 2 surviving recombinant strains was verified by Southern analysis of the chromosomal DNA. High molecular weight DNA was isolated, digested with BamHI and KpnI and subsequently separated by electrophoresis on a 0.7% agarose gel. Following transfer to nitrocellulose hybridization was performed according to standard procedures using the probes described in the previous section. A schematic presentation of the hybridizing fragments is shown in Figure 11C. The results of the Southern analyses are presented in Figure 10B. The presence of a 5.2 kb hybridizing BamHI fragment and a 3.4 kb hybridizing KpnI fragment, with the concomitant loss of the 5.5 kb BamHI and the 6.3 kb hybridizing KpnI fragments is specific for the absence of the amdS selection marker. The weaker hybridizing 7 and 12 kb fragments in a BamHI digest and the 4, 8 and 15 kb KpnI fragments again refer to the endogenous xylanase locus. Both strains show the expected pattern. In these recombinant strains, which were designated GBA-107 and GBA-108, the preferred glaA sequences are removed correctly and that possess finally no selection marker gene at all. Both strains can be reused again to delete or insert other genes or DNA elements by using the same type of vector.

Example 2

Marker gene free introduction of the glaA gene targeted at the 3'glaA non-coding region of the truncated glaA locus in A.niger GBA-107

[0119] In this example the introduction of a gene into the genome of A.niger is described by using approximately the same approach and procedures as described in the previous example. Besides the desired gene or DNA element the vector contains DNA sequences homologous to the host genome to target the vector at a predefined genomic locus of the host, by a single cross-over event. This type of vector comprises a selection marker gene flanked by DNA repeats as well. The selection marker gene in transformants derived with this vector can be removed properly again by applying the counter-selection procedure. As an example the introduction of a glaA gene copy is described which becomes integrated at the truncated glaA locus in the recombinant Δ glaA A.niger GBA-107 strain derived in Example 1 (for a schematic drawing see Figure 12)

Description of the glaA integration vector: pGBGLA30

[0120] The integration vector pGBGLA30 consists of the A.niger amyloglucosidase (glaA) gene under control of the native promoter and the A.nidulans amdS gene under control of the A.nidulans gpdA promoter flanked by 3'glaA non-coding sequences to direct integration at the 3' glaA non-coding region and to remove the amdS selection marker gene via the counter-selection.

Construction of the integration vector

[0121] A 1.8 kb XhoI/EcoRI glaA promoter fragment from pAB6-1 (Figure 13) was subcloned into the SmaI and EcoRI sites of pTZ19R (United States Biochemicals). The protruding 5' end of the XhoI site of the glaA promoter fragment was filled in using the Klenow fragment of E.coli DNA polymerase I prior to cloning in pTZ19R. The SmaI site is destroyed and the XhoI site is restored by this cloning procedure. The thus obtained plasmid was designated pGBGLA5 (Figure 13).

[0122] To introduce appropriate restriction sites (AatII, SnaBI, AsnI and NotI) and to destroy the XhoI site in the glaA promoter, the synthetic fragment consisting of the two oligonucleotides AB3657 (SEQ ID NO: 10) and AB3658 (SEQ ID NO: 11):

```

5' AGCTTGACGTCTACGTATTAATGCGGCCGCT      3' AB3657
      |||||
3'      ACTGCAGATGCATAATTACGCCGCGCAAGCT    5' AB3658

```

was inserted into the HindIII and XhoI sites of pGBGLA5. The thus obtained plasmid was designated pGBGLA26

(Figure 14).

[0123] Next, the 3.4 kb EcoRI fragment from pAB6-1 containing the remaining 3' part of the glaA promoter, the glaA coding sequence and part of the 3' glaA non-coding sequence, was cloned into the EcoRI site of pGBGLA26. This new plasmid was designated pGBGLA27 (Figure 15). This plasmid was partially digested with EcoRI and the synthetic fragment consisting of the oligonucleotides AB3779 (SEQ ID NO: 12) and AB3780 (SEQ ID NO: 13):

5' AATTGGGGCCCATTAAC TCGAGC 3' AB3779
|||||
3' CCCCCGGGTAATTGAGCTCGTTAA 5' AB3780

was inserted into the EcoRI site at the end of the 3' glaA non-coding sequence from the glaA gene. By this cloning step, the EcoRI site was destroyed and an Apal and XhoI restriction site were introduced. The resultant plasmid was designated pGBGLA42 (Figure 16).

[0124] Amplification of the 2.2 kb 3' glaA non-coding sequences and concomitant adjustment of appropriate restriction sites was performed by the Polymerase Chain Reaction (PCR) method.

[0125] In these PCR reactions, plasmid pAB6-1 containing the entire *glaA* locus was used as template and as primers four synthetic oligo nucleotides were designed possessing the following sequence:

Oligo AB3448 (SEQ ID NO: 14):

5' GTGCGAGGTACCACAATCAATCCATTTCGC 3'

(a 3' glaA non-coding specific sequence just downstream the stopcodon of the glaA gene)

Oligo AB3449 (SEQ ID NO: 15):

5' ATGGTTCAAGAACTCGGTAGCCTTTTCCTTGATTCT 3'

(a 3' glaA non-coding specific sequence around the KpnI site approx. 1 kb downstream of the stop codon)

Oligo AB3450 (SEQ ID NO: 16):

5' AGAATCAAGGAAAAGGCTACCGAGTTCTTGAACCAT 3'

(a 3' glaA non-coding specific sequence around the KpnI site approx. 1 kb downstream of the stop codon)

Oligo AB3520 (SEQ ID NO: 17):

5' ATCAATCAGAAGCTTTCTCTCGAGACGGGCATCGGAGTCCCCG 3'

(a 3' glaA non-coding specific sequence approx. 2.2 kb downstream of the stopcodon)

[0126] To destroy the KpnI site approximately 1 kb downstream of the stop codon from the glaA gene and to alter the SaiI site approximately 2.2 kb downstream the stop codon from the glaA gene into a XhoI site two separate polymerase chain reactions were performed: the first reaction with oligonucleotides AB3448 (SEQ ID NO: 14) and AB3449 (SEQ ID NO: 15) as primers to amplify an approximately 1 kb DNA fragment just downstream the stopcodon of the glaA gene, and the second reaction with oligonucleotides AB3450 (SEQ ID NO: 16) and AB3520 (SEQ ID NO: 17) as primers to amplify an approximately 1.2 kb DNA fragment just downstream the KpnI site in the 3' glaA non-coding region both with pABG-1 as template. A schematic view of these amplifications is presented in Figure 17A. The PCR

was performed as described in example 1. Twenty-five amplification cycles (each 1 minute at 55°C; 1.5 minutes at 72°C and 1 minute at 94°C) were carried out.

[0127] The two generated PCR DNA fragments were purified by agarose gel electrophoresis and ethanol precipitation and subsequently used as template in the third PCR with oligonucleotides AB3448 (SEQ ID NO: 14) and AB3520 (SEQ ID NO: 17) as primers to generate the fusion fragment. Twenty-five amplification cycles (each: 2 minutes at 55°C; 3 minutes at 72°C; 2 minutes at 94°C) were carried out in a DNA-amplifier (Perkin-Elmer/Cetus). The amplified DNA fragment was purified by agarose gel electrophoresis and ethanol precipitation and subsequently subcloned in the *Sma*I site of pTZ18R. The obtained plasmid was designated pGBGLA17 (Figure 17B).

[0128] To fuse this adjusted 3' *glaA* non-coding region to the *amdS* gene, a part of the *amdS* gene was subcloned from pGBDEL4L into pSP73 (Promega). For this construction, pGBDEL4L was digested with *Bgl*III and *Hind*III, the 3.4 kb *amdS*/3' *glaA* non-coding fragment was isolated by agarose gel electrophoresis and subcloned into the appropriate sites of pSP73 (Promega). The resulting plasmid was designated pGBGLA21 (Figure 18).

[0129] The approximately 1 kb sized 3' *glaA* non-coding region in this plasmid was exchanged by the 2.2 kb 3' *glaA* non-coding region of pGBGLA17. pGBGLA17 and pGBGLA21 were digested with *Kpn*I and *Hind*III. The 2.2 kb 3' *glaA* non-coding region DNA fragment from pGBGLA17 and the 4.9 kb DNA fragment of pGBGLA21 were isolated by agarose gel electrophoresis, ligated and subsequently molecular cloned by transferring the ligation mixture to *E. coli*. The thus derived plasmid was designated pGBGLA22 (Figure 19).

[0130] The *amdS* gene with the extended 3' *glaA* non-coding region was completed with the *gpdA* promoter and fused to the remaining part of the *amdS* gene. pGBGLA22 was digested with *Bgl*III and *Hind*III, the 4.4 kb *amdS*/3' *glaA* non-coding region DNA fragment isolated by agarose gel electrophoresis, subsequently ligated with plasmid pGBGLA24 digested with *Bgl*III and *Hind*III and transferred to *E. coli*. The thus derived plasmid was designated pGBGLA25 (Figure 20).

[0131] pGBGLA25 was partially digested with *Eco*RI and in the *Eco*RI site of the *gpdA* promoter the synthetic fragment consisting of the two oligonucleotides AB3781 (SEQ ID NO: 18) and AB3782 (SEQ ID NO: 19):

```

5'   AATTGGGGCCCGAGCGTCC           3'   AB3781
      |||||
3'   CCCCGGGTCGCGAGGTTAA           5'   AB3782

```

was inserted. This new plasmid was designated pGBGLA43 (Figure 21). Due to this cloning step, the *Eco*RI restriction site just in front of the *gpdA* promoter was destroyed by the introduction of an *Apa*I restriction site.

[0132] The plasmid pGBGLA43 was digested with *Apa*I and *Xho*I, and the 5.3 kb DNA fragment comprising the *gpdA* promoter/*amdS* gene/3' *glaA* non-coding region was isolated by agarose gel electrophoresis, subsequently ligated with pGBGLA42 digested with *Apa*I and *Xho*I, and transferred to *E. coli*. The derived plasmid was designated pGBGLA28 (Figure 22).

[0133] Prior to cloning, the 3' *glaA* non-coding region DNA fragment (positioned at approximately 2.2 kb downstream the stop codon of the *glaA* gene, designated 3' *glaA* non-coding region), was amplified and provided with suitable restriction sites using the PCR method.

[0134] For this PCR reaction, the plasmid pAB6-1 was used as template and as primers two synthetic oligonucleotides were designed possessing the following sequence:

Oligo AB3746 (SEQ ID NO: 20):

```
5' TGACCAATAAAGCTTCTCGAGTAGCAAGAAGACCCAGTCAATC 3'
```

(a partly 3' *glaA* non-coding specific sequence around the *Sal*I site positioned at about 2.2 kb downstream the stop codon of the *glaA* gene).

Oligo AB3747 (SEQ ID NO: 21):

5'CTACAAACGGCCACGCTGGAGATCCGCCGGCGTTCGAAATAACCACT3'

(a partly 3''glaA non-coding specific sequence around the XhoI site located at about 4.4 kb downstream the stop codon of the glaA gene)

[0135] Twenty-five amplification cycles (each: 1 minute 55°C; 1.5 minutes 72°C; 1 minute 94°C) were carried out in a DNA-amplifier (Perkin-Elmer/Cetus). A schematic representation of this amplification is shown in figure 23A. The thus obtained DNA fragment was digested with HindIII, purified by agarose gel electrophoresis and ethanol precipitation and subcloned in both orientations into the HindIII site of pTZ19R. The resulting plasmids were designated pGBGLA29A and pGBGLA29B (Figure 23).

[0136] The final step comprises the insertion of the 3'glaA non-coding sequence from pGBGLA29A into the plasmid pGBGLA28. To achieve this, pGBGLA29A was digested with HindIII and NotI. The 2.2 kb sized 3'glaA non-coding region fragment was isolated by agarose gel electrophoresis, subsequently ligated to pGBGLA28 digested with HindIII and NotI and transferred to E. coli. The derived integration vector was designated pGBGLA30 (Figure 24).

Transformation of A.niger GBA-107 with the integration vector pGBGLA30

[0137] Prior to transformation, E.coli sequences were removed from the integration vector pGBGLA30 by XhoI digestion and agarose gel electrophoresis. The A.niger strain GBA-107 was transformed with either 5 or 10 µg DNA fragment by procedures as described in the experimental section. Single A.niger transformants were purified several times on selective acetamide containing plates. Spores of individual transformants were collected following growth for about 5 days at 30°C on 0.4% potato dextrose agar (Oxoid, England) plates. Southern analyses were performed to verify whether integration into the 3' glaA non coding region of the endogenous truncated glaA locus had occurred. High molecular weight DNA of several transformants was isolated, digested with either KpnI, or BglII and subsequently fractionated by electrophoresis on a 0.7% agarose gel. After transfer to nitrocellulose filters, hybridization was performed according to standard procedures. As probe a ³²P-labelled approx. 0.7 kb XhoI/SalI glaA promoter fragment isolated from plasmid pAB6-4 (described in example 1) was used. The results of only 3 transformants (#107-5, #107-9 and #107-7) and the reference strain A.niger GBA107 and its ancestor A.niger CBS 531.88 are shown as example in Figure 25. For a better understanding of the autoradiograph, a schematic presentation is given in Figure 26A,B,C showing the sizes of the hybridizing fragments of the intact glaA locus, the truncated glaA locus and of the truncated glaA locus with a single pGBGLA30 copy integrated into the predefined 3' glaA non-coding region.

[0138] Characteristic for the intact glaA locus is a 4.5 kb hybridizing fragment in a KpnI digest and a 10 kb hybridizing fragment in a BglII digest. Characteristic for the truncated glaA locus of A.niger GBA-107 is a 3.4 kb hybridizing fragment in a KpnI digest and a 13 kb hybridizing fragment in a BglII digest. In case of integration of the pGBGLA30 vector into the 3' region of the truncated glaA locus, in a KpnI digest an additional 6.7 kb hybridizing fragment is expected besides the 3.4 kb hybridizing fragment and in a BglII digest the 13 kb hybridizing fragment is absent and replaced by a 14.5 kb hybridizing fragment. As can be seen in Figure 25, transformants #107-5 and #107-9 show the expected hybridization pattern of a single pGBGLA30 copy integrated into the predefined 3' non-coding region of the truncated glaA locus. The hybridization pattern of transformant #107-7 indicates integration of the pGBGLA30 copy elsewhere into the genome of A.niger GBA-107. The transformants with the correctly integrated pGBGLA30 copy were designated GBA-119 and GBA-122 and were used to remove subsequently the amdS selection marker gene properly.

Removal of the amdS selection marker gene from A.niger GBA-119 and GBA-122 by counter-selection on fluoracetamide containing plates.

[0139] The amdS selection marker gene in the transformants A.niger GBA-119 and GBA-122 was removed again as described in the experimental section. The removal of the amdS selection marker gene in several surviving recombinant strains was verified by Southern analysis of the chromosomal DNA. High molecular weight DNA was isolated, digested either with KpnI or BglII and subsequently separated by electrophoresis on a 0.7% agarose gel. Following transfer to nitrocellulose, hybridization was performed according to standard procedures. As probe the ³²P labelled 2.2 kb HindIII/NotI 3'glaA non-coding fragment isolated from plasmid pGBGLA29A (described previously, Figure 24) was used.

[0140] A schematic presentation of the hybridizing fragments is shown in Figure 26. The results of only 3 surviving recombinant strains from A.niger GBA-119 (#AG5-5, #AG5-6 and #AG5-7) as well as 3 surviving recombinant strains

from *A.niger* GBA-122 (#AG9-1, #AG9-2 and #AG9-4) and the reference strains *A.niger* CBS 531.88 and *A.niger* GBA-107 are shown in Figure 27A,B.

[0141] In strain *A.niger* CBS 531.88 a 6.9 kb hybridizing fragment is present in a *KpnI* digest and a 6.9 kb hybridizing fragment in a *BglII* digest. In the *A.niger* GBA-107 strain a 6.9 kb hybridizing fragment is present in a *KpnI* digest and a 13 kb hybridizing fragment in a *BglII* digest. In the *A.niger* strains GBA-119 and GBA-122 with a single pGBGLA30 copy integrated into the 3' *glaA* non-coding region an 8 kb and a 6.7 kb hybridizing band are present in a *KpnI* digest and a 14.5 kb and a 7.6 kb hybridizing band are present in a *BglII* digest.

[0142] Specific for correct removal of the *amdS* selection marker gene is the presence of a 6.7 kb and a 8.5 kb hybridizing fragment in a *KpnI* digest and concomitant loss of the 8 kb hybridizing fragment. In a *BglII* digest, a 14.5 kb and a 6.9 kb hybridizing fragment with concomitant loss of the 7.6 kb hybridizing fragment is specific for the absence of the *amdS* selection marker gene. As can be seen in Figure 27, strains #AG5-7, #AG5-5, #AG9-1 and #AG9-4 show the expected hybridizing pattern of the correctly removed *amdS* selection marker gene. These strains were designated GBA-120, GBA-121, GBA-123 and GBA-124 respectively. The hybridizing patterns of strains #AG5-6 and #AG9-2 indicate loss of the entire pGBGLA30 copy resulting in the parental *A.niger* GBA-107 strain with only a truncated *glaA* locus.

[0143] Strains *A.niger* GBA-120, GBA-121, GBA-123 and GBA-124 were tested in shake flask fermentations for the ability to produce glucoamylase. As reference strains *A.niger* CBS 531.88, GBA-107, GBA-119 and GBA-122 were tested. Shake flask fermentations and the glucoamylase assay were performed as described in the experimental section. In the strains GBA-119 till GBA-124 levels varying between 150-200 U/ml could be measured. These glucoamylase levels were to be expected and comparable to levels obtained with the parental untransformed wild-type strain *A.niger* CBS 531.88.

Example 3

Marker gene free introduction of the phytase gene targeted at the 3' *glaA* non-coding region of the truncated *glaA* locus in *A.niger* GBA-107

[0144] In this example describes the introduction of a gene into the genome of *A.niger* by using approximately the same approach and procedures as described in the previous example. The main difference is that the gene of interest and the selection marker gene are located on two separate vectors and that these vectors are co-transformed to *A.niger*. Besides the gene of interest or the marker gene, the vectors contain DNA sequences homologous to the host genome to target the vectors at a predefined genomic locus of the host, by a single cross-over event. By performing the fluoracetamide counter-selection on these (co)-transformants (as described in the experimental procedures), the *amdS* marker gene will be deleted properly by an internal recombination event between the DNA repeats that are created by integration via a single cross-over event.

Description of the vectors used for co-transformation

[0145] The vector with the gene of interest pGBGLA53 consists of the *A.ficuum* phytase gene under control of the *A.niger* glucoamylase (*glaA*) promoter flanked by 3' *glaA* non-coding sequences to direct integration at the 3' *glaA* non-coding region. The vector with the selection marker gene pGBGLA50 consists of the *A.nidulans* *amdS* gene under control of the *A.nidulans* *gpdA* promoter flanked by 3' *glaA* non-coding sequences to direct integration at the 3' *glaA* non-coding region.

Construction pathway of pGBGLA50

[0146] The construction of pGBGLA50 comprises one cloning step. Plasmid pGBGLA29A was digested with *HindIII* and the sticky ends were filled in using the Klenow fragment of *E.coli* DNA polymerase. Next, the 2.2 kb 3' *glaA* non-coding region fragment was isolated by agarose gel-electrophoresis, subsequently ligated into pGBGLA43 digested with *ApaI* and treated with T4 DNA polymerase to generate blunt ends, and transferred to *E.coli*. The derived plasmid with the 3' *glaA* non-coding region DNA fragment in the correct orientation was designated pGBGLA50 (Figure 28).

Construction pathway of pGBGLA53

[0147] The first step in the construction pathway of pGBGLA53 is the subcloning of two fragments, comprising the *glaA* promoter fused to almost entire coding sequence of the *A.ficuum* phytase gene. To achieve this, plasmid pGBGLA42 was digested with *HindIII* and *EcoRI* and the 1.8 kb *HindIII/EcoRI* 5' *glaA* promoter fragment was isolated by agarose gel-electrophoresis. Plasmid pFYT3 (European Patent Application 0 420 358 A1) was digested with *EcoRI*

and BglII and the 1.6 kb EcoRI/BglII fragment comprising the 3'part of the glaA promoter fused to the 5' part of the phytase gene was isolated by agarose gel-electrophoresis and ligated together with the 1.8 kb HindIII/EcoRI 5'glaA promoter fragment isolated from pGBGLA42 into the HindIII and BglII sites of pSp73 (Promega). The resulting plasmid was designated pGBGLA49 (Figure 29).

[0148] The next step is the cloning of a 3'glaA non-coding region DNA fragment into pGBGLA49. Prior to cloning, this 3'glaA non-coding region DNA fragment (positioned at approximately 2.2 kb downstream the stop codon of the glaA gene) was amplified and provided with suitable restriction sites using the PCR method.

[0149] For this PCR reaction, the plasmid pAB6-1 was used as template and as primers two synthetic oligonucleotides with the following sequence were designed:

Oligo AB4234 (SEQ ID NO: 22):

5' GAAGACCCAGTCAAGCTTGCATGAGC 3'

(a 3'glaA non-coding sequence located approximately 2.2 kb downstream the stopcodon of the glaA gene)

Oligo AB 4235 (SEQ ID NO: 23):

5'TGACCAATTAAGCTTGC GGCCGCTCGAGGTCGCACCGGCAAAC 3'

(a 3'glaA non-coding sequence located approximately 4.4 kb downstream the stopcodon of the glaA gene)

[0150] Twenty-five amplification cycles (each: 1 minute 94°C; 1 minute 55°C; 1.5 minutes 72°C) were carried out in a DNA-amplifier (Perkin-Elmer). A schematic representation of this amplification is shown in figure 30A. The thus obtained fragment was digested with HindIII, purified by agarose gel-electrophoresis and subcloned into the HindIII site of pTZ19R. The resulting plasmid was designated pGBGLA47 (Figure 30).

[0151] Plasmid pGBGLA47 was digested with HindIII en NotI, the 2.2 kb 3'glaA non-coding DNA fragment was isolated by agarose gel-electrophoresis and cloned into the HindIII and NotI sites of pGBGLA49. The resulting plasmid was designated PGBGLA51 (Figure 31).

[0152] The last step in the construction pathway of pGBGLA53 is the cloning of the DNA fragment comprising the remaining part of the phytase coding sequence fused to the 3'glaA non-coding DNA fragment located just downstream the stop codon of the glaA gene. Prior to cloning, the remaining part of the phytase gene and the 3'glaA non-coding DNA fragment located just downstream the stopcodon of the glaA gene were fused and provided with suitable restriction sites using the PCR method. In the PCR, plasmid pAB6-1 was used as template and as primers two synthetic oligonucleotides were used, having the following sequences:

Oligo AB4236 (SEQ ID NO: 24):

5' TGACCAATAAAGCTTAGATCTGGGGGTGATTGGGCGGAGTGTGCTT
AGACAATCAATCCATTTTCGC 3'

(36 bp of the phytase coding sequence, starting at the BglII site until the stopcodon fused to the 3'glaA non-coding region, starting just downstream the stopcodon of the glaA gene)

Oligo AB4233 (SEQ ID NO: 25):

5' TGACCAATAGATCTAAGCTTGACTGGGTCTTCTTGC 3'

(a 3' glaA non-coding sequence located approximately 2.2 kb downstream the stopcodon of the glaA gene)

[0153] Twenty-five amplification cycles (each: 1 minute 94°C; 1 minute 55°C; 1.5 minutes 72°C) were carried out in a DNA-amplifier (Perkin-Elmer). A schematic representation of this amplification is shown in figure 32A. The thus obtained fragment was digested with HindIII, purified by agarose gel-electrophoresis and subcloned in both orientations into the HindIII site of pTZ19R. The resulting plasmids were designated pGBGLA48 and pGBGLA52 (figure 32B).

[0154] Plasmid pGBGLA52 was digested with BglII and partially digested with BamHI, the 2.2 kb phytase/3'glaA non-coding DNA fragment was isolated by agarose gel-electrophoresis and cloned into the BglII site of pGBGLA51. The derived plasmid with the 2.2 kb phytase/3'glaA non-coding DNA fragment in the correct orientation was designated pGBGLA53 (figure 33).

Transformation of A.niger GBA-107 with the vectors pGBGLA50 and pGBGLA53

[0155] Prior to transformation, E.coli sequences were removed from pGBGLA50 and pGBGLA53 by respectively XhoI or HindIII digestion followed by agarose gel-electrophoresis. The A.niger GBA-107 strain was transformed with respectively 1 µg pGBGLA50 fragment plus 1 µg pGBGLA53 fragment, 1 µg pGBGLA50 fragment plus 5 µg pGBGLA53 fragment, or 1 µg pGBGLA50 fragment plus 10 µg pGBGLA53 fragment using the transformation procedure described in the experimental section.

[0156] Single transformants were isolated, purified and Southern analysis was performed, using the same digests and probes as described in example 2, to verify integration of both pGBGLA50 and pGBGLA53. In about 10-20% of the analyzed transformants both pGBGLA50 and pGBGLA53 were integrated into the genome of the A.niger GBA-107 host strain. The transformant showing the correct integration pattern of a single copy pGBGLA50 and a single copy pGBGLA53, both integrated at the predefined 3'glaA non-coding region of the truncated glaA locus was used to remove subsequently the amdS selection marker gene.

Removal of the amdS marker gene by counter-selection on fluoracetamide containing plates

[0157] By performing the fluoracetamide counter-selection (as described in the experimental procedures), the amdS marker gene was deleted by an internal recombination event between the DNA repeats that were created by integration via a single cross-over event (i.e. the 3'glaA non-coding sequences). Proper removal of only the amdS marker gene was verified by Southern analysis using the same digests and probes as in example 2.

Example 4

Marker gene free introduction of the glaA gene and the phytase gene in A.oryzae

[0158] This example describes the marker gene free introduction of the glaA gene or the phytase gene in A.oryzae NRRL3485. A.oryzae NRRL3485 was transformed as described in the experimental section using the same vectors and approach as described in examples 2 and 3. Single transformants were isolated, purified and Southern analysis of chromosomal DNA of several transformants was performed to verify integrations of respectively the pGBGLA30 vector or the pGBGLA50 and pGBGLA53 vectors. In the Southern analysis, the same digests and probes were used as described in example 2.

Removal of the amdS gene by counter-selection on fluoracetamide containing plates

[0159] In case of integration of the pGBGLA30 vector, a transformant with a single copy of the pGBGLA30 integrated into the genome of the host strain A.oryzae NRRL3485 was used to remove the amdS gene properly. The counter-selection on fluoracetamide containing plates was performed as described in the experimental section. Correct removal of the amdS gene was verified by Southern analysis of chromosomal DNA of several fluoracetamide resistant strains. The same digests and probes were used as described in Example 2.

[0160] In case of co-transformation of the pGBGLA50 and pGBGLA53 vector, a transformant with a single copy of

both pGBGLA50 and pGBGLA53 integrated into the host genome was used to remove the amdS marker gene properly. The counter-selection using fluoracetamide plates was performed as described in the experimental section. Correct removal of the amdS marker gene (e.g. the pGBGLA50 vector) was verified by Southern analysis of chromosomal DNA of several fluoracetamide resistant strains using the same digests and probes as described in example 2.

Example 5

Marker gene free introduction of the glaA gene and the phytase gene in T.reesei

[0161] This example describes the marker gene free introduction of the glaA gene or the phytase gene in Trichoderma reesei strain QM9414 (ATCC 26921). T.reesei QM9414 was transformed as described in the experimental section using the same vectors and approach as described in examples 2 and 3. Single transformants were isolated, purified and Southern analysis of chromosomal DNA of several transformants was performed to verify whether integration of respectively the pGBGLA30 vector or the pGBGLA50 and pGBGLA53 vectors. In the Southern analysis, the same digests and probes were used as described in example 2.

Removal of the amdS gene by counter-selection on fluoracetamide containing plates

[0162] In case of integration of the pGBGLA30 vector, a transformant with a single copy of the pGBGLA30 integrated into the genome of the host strain T.reesei QM9414 was used to remove the amdS gene properly. The counter-selection on fluoracetamide containing plates was performed as described in the experimental section. Correct removal of the amdS gene was verified by Southern analysis of chromosomal DNA of several fluoracetamide resistant strains.

[0163] In case of co-transformation of the pGBGLA50 and pGBGLA53 vector, a transformant with a single copy of both pGBGLA50 and pGBGLA53 integrated into the host genome was used to remove the amdS marker gene properly. The counter-selection using fluoracetamide plates was performed as described in the experimental section. Correct removal of the amdS marker gene (e.g. the pGBGLA50 vector) was verified by Southern analysis on chromosomal DNA of several fluoracetamide resistant strains using the same digests and probes as described in example 2.

Example 6

Marker gene free introduction into P.chrysogenum of a P.chrysogenum gene by co-transformation using the amdS-gene as a selection marker

[0164] In this example the marker gene free introduction of a gene into the genome of P.chrysogenum by co-transformation is described.

[0165] In the co-transformation procedure, 2 different pieces of DNA are offered to the protoplasts, one of them being the amdS-selection marker, on the presence of which the first transformant selection takes place, as described in the experimental section, the second being another piece of DNA of interest, e.g. encoding a particular enzyme of interest. In a certain number of transformants both pieces of DNA will integrate into the chromosomes and will be stably maintained and expressed.

[0166] The amdS-selection marker gene can then be removed selectively from purified transformants by applying the counter-selection procedure as described in the experimental section, while the second piece of DNA will remain stably integrated into the chromosomes of the transformant. As an example to illustrate the general applicability of the method the stable, marker gene free introduction of a niaD-gene is described which enables a niaD-host to grow on nitrate as sole nitrogen-source.

[0167] Host for this co-transformation is a P.chrysogenum niaD-strain which lacks nitrate reductase and therefore is unable to grow on plates containing nitrate as sole nitrogen source. These strains can be easily obtained by well known procedures (Gouka et al., Journal of Biotechnology 20(1991), 189-200 and references there in)

[0168] During the co-transformation (procedure described in experimental section), two pieces of DNA are simultaneously offered to the protoplasts: the 7.6 kb EcoRI restriction fragment from pGBGLA28 containing the amdS selection marker gene and the 6.5 kb EcoRI restriction fragment from pPC1-1, containing the P.chrysogenum niaD-gene. Prior to transformation, both fragments have been separated from E.coli vector sequences by agarose gel-electrophoresis and purified from agarose gel by electro-elution. The first selection of transformants took place on selective plates containing acetamide as sole nitrogen source as described in the experimental section.

[0169] Among the transformants, co-transformants are found by replica plating spores of purified transformants to plates containing nitrate as sole nitrogen source. Typically about 20-60% of the replica plated transformants were able to grow on this medium, indicating that in these transformants not only the amdS selection marker gene but also the niaD-gene has integrated into the genome and is expressed.

Removal of the amdS gene by counter-selection on fluoracetamide containing plates

[0170] The amdS selection marker gene is subsequently removed from the co-transformants by counter-selection on fluoracetamide.

5 [0171] For direct selection on the amdS⁻/niaD⁺-phenotype the medium used contained 10 mM fluor-acetamide. Spores were plated at a density of 10⁴ spores per plate. After 5-7 days of incubation at 25°C, fluor-acetamide resistant colonies could be identified as solid colonies clearly distinct from the faint background. The niaD⁺-phenotype of the recombinants is demonstrated by their growth on the fluoracetamide-medium containing nitrate as sole nitrogen source. The amdS⁻-phenotype of the recombinants was confirmed by lack of growth of the recombinants on plates containing acetamide as sole nitrogen source. Typically, 0.1-2% of the original number of plated spores exhibited the desired phenotype.

[0172] Southern analysis on chromosomal DNA from several fluoracetamide resistant strains confirmed that the amdS selection marker gene was removed from the P.chrysogenum genome.

15 Example 7

Test of the amdS-minus phenotype of the yeast Kluyveromyces lactis

20 [0173] A prerequisite for the use of the amdS selection system in K.lactis is that this yeast does not contain any acetamidase activity. To test this we have plated the K.lactis strains CBS 683 and CBS 2360 on the following 3 different solid media:

- I Yeast Carbon Base (YCB, Difco), containing all the essential nutritives and vitamins except a nitrogen-source.
- II YCB supplemented with 5mM acetamide.
- 25 III YCB supplemented with 0.1 % (w/v) NH₄(SO₄)₂.

All 3 media contained 1.2 % (w/v) Oxoid agar (Agar No. 1) and 30 mM Sodium Phosphate buffer at pH 7.0. Difco YCB was used at 1.17 % (w/v).

30 [0174] Full grown K.lactis colonies were only observed on medium III, containing ammonium as nitrogen source. In plates without nitrogen-source or with acetamide as sole nitrogen-source no growth or, occasionally slight background growth was observed, which is most likely caused by trace amounts of nitrogen contaminating the agar or other medium components. We conclude that both K.lactis strains lack sufficient acetamidase activity to sustain growth on acetamide as sole nitrogen source. This should allow for the A.nidulans amdS gene to be used as selection marker in the yeast K.lactis.

35 Example 8

Construction of plasmids for use of the amdS gene in yeasts

40 Construction of pGBamdS1

[0175] We have previously used pGBHSA20 for the expression of human serum albumin (HSA) in K.lactis (Swinkels et al. 1993, Antonie van Leeuwenhoek 64, 187-201). In pGBHSA20 the HSA cDNA is driven from the K.lactis LAC4 promoter (Figure 34 for the physical map of the plasmid pGBHSA20). At the 3'-end the HSA cDNA is flanked by LAC4 terminator sequences. For selection of transformants pGBHSA20 contains the Tn5 phosphotransferase gene which confers resistance to the antibiotic G418 (Geneticin, BRL) (Reiss et al. (1984) EMBO J. 3, 3317-3322) driven by the S.cerevisiae ADH1 promoter (Bennetzen and Hall (1982) J. Biol. Chem. 257, 3018-3025). In the unique SstII site of the LAC4 promoter pGBHSA20 contains the E.coli vector pTZ19R which is used for amplification in E.coli. Prior to transformation to K.lactis the pTZ19R sequences are removed from pGBHSA20 by SstII digestion and agarose gel purification. Transformation of pGBHSA20 linearized in the SstII site of the LAC4 promoter to K.lactis results in integration into the genomic LAC4 promoter by homologous recombination. pGBamdS1 is derived from pGBHSA20 by substitution of the HSA cDNA for the amdS cDNA from pamdS1. Using PCR, SaiI sites were introduced at the 5' and 3' ends of the amdS cDNA. In this PCR pamdS1 was used as template and oligo's AB3514 (SEQ ID NO: 26) and AB3515 (SEQ ID NO: 27) were used as primers.

55

Oligo AB3514 (SEQ ID NO: 26):

5'-CTGCCGAATTCGTCGACATGCCTCAATCCTGGG-3'

(an 5' end amdS-specific sequence with the introduced SalI site)

Oligo AB3515 (SEQ ID NO: 27):

5'-GGCAGTCTAGAGTCGACCTATGGAGTCACCACATTTTC-3'

(an 3' end amdS-specific sequence with the introduced SalI site).

[0176] The PCR fragment thus obtained was digested with SalI and cloned into the SalI/XhoI sites of pGBHSA20. Several clones were obtained containing either of the 2 possible orientations of the amdS cDNA as judged by restriction analysis. One of the clones with the amdS cDNA in the correct orientation is pGBamdS1 the physical map of which is shown in Figure 34.

Construction of pGBamdS3

[0177] By heterologous hybridization using a probe derived from the S.cerevisiae elongation factor 1- α gene (EF1- α ; Nagata et al. (1984) EMBO J. 3, 1825-1830), we have isolated a genomic clone containing the K.lactis homologue of the EF1- α gene, which we call KIEF1. In this example we have used a 813 bp fragment containing the KIEF1 promoter to express the amdS cDNA in K.lactis: Using oligonucleotides AB3701 (SEQ ID NO: 28) and AB3700 (SEQ ID NO: 29), this fragment was amplified in a PCR using genomic DNA from K.lactis strain CBS 683 as template. AB3700 (SEQ ID NO: 29) is designed such that it contains 21 nucleotides of the KIEF promoter and 38 nucleotides upstream the ATG initiation codon of the amdS gene. The sequence of AB3701 (SEQ ID NO: 28) and AB3700 (SEQ ID NO: 29) is as shown:

Oligo AB3701 (SEQ ID NO: 28):

5'-CTGCCGAATTCGTCGACACTAGTGGTACCATTATAGCCATAGGACAGCAAG 3'

(a 5' KIEF1-specific promoter sequence with the additional restriction sites EcoRI, SalI, SpeI and KpnI at the 5' end of the promoter)

Oligo AB3700 (SEQ ID NO: 29):

5'-GCTCTAGAGCGCGCTTATCAGCTTCCAGTTCTTCCCAGGATT-GAGGCATTTTTTAATGTTACTTCTCTTGC-3'

(3' KIEF1-specific promoter sequence fused to the 5'-sequences of the amdS cDNA with the restriction sites BssH2 and additional site XbaI).

[0178] The PCR was performed using standard conditions and the PCR-fragment obtained was digested with EcoRI and XbaI and subcloned into EcoRI/XbaI digested pTZ19R. The physical map of the resulting plasmid pTZKIEF1 is shown in Figure 35. The remaining part of the amdS cDNA as well as part of the LAC4 terminator sequences were obtained from pGBamdS1 by digestion with BssH2 and SphI. This BssH2-SphI fragment was cloned into the BssH2 and SphI digested pTZKIEF1 and the resulting plasmid was designated pGBamdS2 (Figure 35). For the final step in

the construction of pGBamdS3, both pGBamdS2 and pTY75LAC4 (Das and Hollenberg (1982) Current Genetics 6, 123-128) were digested with *Sph*I and *Hind*III. The 5.7 kb DNA fragment from pGBamdS2 and the 1.2 kb DNA fragment from pTY75LAC4, which contains the remaining *LAC4* terminator sequences, were purified from agarose gels after fractionation and subsequently ligated and used to transform *E.coli*. The resulting expression vector, in which the *amdS* cDNA is driven from the *K.lactis* *KIEF1* promoter, was designated pGBamdS3 (Figure 36).

Construction of pGBamdS5

[0179] Fusion of the *S.cerevisiae* alcohol dehydrogenase I (*ADH1*) promoter to the *amdS* cDNA was performed in a PCR using pGBHSA20 as template. One of the primers (AB3703; SEQ ID NO: 31) contains sequences complimentary to the 3'-end of the *ADH1*-promoter sequence which are fused to sequences of the *amdS* cDNA. The other primer (AB3702; SEQ ID NO: 30) contains the 5'-end of the *ADH1* promoter:

Oligo AB3702 (SEQ ID NO: 30):

5'-CTGCGAATTCGTCGACACTAGTGGTACCATCCTTTTGTGTTTCCGGGTG-3'

(a 5' *ADH1*-specific promoter sequence with the additional restriction sites *Eco*RI, *Sal*I, *Spe*I and *Kpn*I at the 5' end of the promoter).

Oligo AB3703 (SEQ ID NO: 31):

5'-GCTCTAGAGCGCGCTTATCAGCGGCCAGTTCTTCCCAGGAT-TGAGGCATTGTATATGAGATAGTTGATTG-3'

(a 3' *ADH1*-specific promoter sequence fused to the 5' *amdS* sequence with additional restriction sites *Bss*H2 and *Xba*I).

[0180] The PCR reaction was performed using a "touchdown" protocol (Don et al., (1991) Nucleic Acids Res. 19, 4008). The reaction mixtures were subjected to 30 amplification cycles, while the annealing temperature was decreased 1°C every two cycles, starting 55°C down to a "touchdown" at 40°C, at which temperature 10 more cycles were carried out (cycles: 2' at 94°C, 2' annealing, 3' at 72°C). The PCR-fragment obtained was digested with *Eco*RI and *Xba*I and subcloned into pTZ19R. The resulting plasmid pTZs.c.ADH1 is shown in Figure 37. pTZs.c.ADH1 and pGBamdS3 were digested with *Kpn*I and *Bss*H2. The 6.8 kb fragment from pGBamdS3 and the 750 bp fragment from pTZs.c.ADH1 were purified by gel electrophoresis, ligated and used to transform *E.coli* JM109. The resulting expression vector was designated pGBamdS5 (Figure 37).

Construction of pGBamdS6

[0181] Plasmid pGBamdS3 contains the *amdS* cDNA under control of the *KIEF1* promoter and flanked at the 3' end by 1.5 kb of *LAC4* terminator sequences (Figure 36). pGBamdS6 is constructed by cloning a fragment which contains a fusion of the *LAC4* promoter and terminator sequences upstream of the *amdS* expression-cassette in pGBamdS3 (Figure 38). In order to fuse the *LAC4* promoter and terminator sequences we have first constructed pPTLAC4 (Figure 39). Using a PCR, additional restriction sites are introduced at the 5' and 3' end of a 600 bp *LAC4* terminator fragment. In the PCR *K.lactis* CBS 683 chromosomal DNA was used as template and oligonucleotides AB3704 (SEQ ID NO: 32) and AB3705 (SEQ ID NO: 33) were used as primers:

Oligo AB3704 (SEQ ID NO: 32):

5'-GCTCTAGAAGTCGACACTAGTCTGCTACGTACTCGAGAATTTATACTTAGA-
TAAG-3'

(a LAC4 terminator-specific sequence starting at the LAC4 stop codon with the additional restriction sites XbaI, SalI, SpeI, SnaBI and XhoI).

Oligo AB3705 (SEQ ID NO: 33):

5'-TGCTCTAGATCTCAAGCCACAATTC-3'

(3' LAC4 terminator-specific sequence with the additional restriction site XbaI).

[0182] The PCR was performed using standard conditions and the resulting DNA fragment was digested with XbaI and subcloned into the XbaI site of pTZ19R to give pTLAC4 (Figure 39). The LAC4 promoter sequence is obtained by digestion of pKS105 van den Berg et al. (1990) Bio/Technology 8, 135-139) with XbaI and SnaBI. The XbaI-SnaBI LAC4 promoter fragment was cloned into the SpeI/SnaBI sites of pTLAC4 and designated pPTLAC4 (Figure 39). For the final step in the construction of pGBamdS6, the plasmid pPTLAC4 was digested with XbaI. The 4.1 kb DNA fragment from pPTLAC4 was purified by gel-electrophoresis and cloned into SpeI site of pGBamdS3. The obtained gene-replacement vector was designated pGBamdS6 (Figure 38).

Construction of pGBamdS8

[0183] pGBamdS7 was constructed by cloning a fragment, which contains part of the LAC4 promoter as well as the chymosin expression-cassette, in between the LAC4 promoter and terminator sequences as present in pGBamdS6 (Figure 40). Plasmid pKS105 contains the prochymosin cDNA fused to the prepro-region of *S.cerevisiae* a-factor under control of the LAC4 promoter (van den Berg et al. (1990) Bio/Technology 8, 135-139). Using a PCR, additional restriction sites were introduced at the 5' and 3' end of the fusion LAC4 promoter and chymosin expression-cassette. In the PCR pKS105 DNA was used as template and oligonucleotides. AB3965 (SEQ ID NO: 34) and AB3966 (SEQ ID NO: 35) were used as primers:

Oligo AB3965 (SEQ ID NO: 34):

5'-CTGCTACGTAATGTTTTTCATTGCTGTTTTAC-3'

(a LAC4 promoter-specific sequence starting at the restriction site SnaBI)

Oligo AB3966 (SEQ ID NO: 35):

5'-CCGCCCAGTCTCGAGTCAGATGGCTTTGGCCAGCCCC-3'

(chymosin-specific sequence with the additional restriction site XhoI).

[0184] The PCR was performed using standard conditions and the obtained PCR fragment was digested with SnaBI and XhoI. The plasmid pGBamdS6 was partially digested with XhoI and subsequently digested with SnaBI and the 10.9 kb DNA fragment was isolated and purified by gel-electrophoresis. The SnaBI-XhoI fusion fragment LAC4 promoter/chymosin expression-cassette was cloned into the SnaBI/XhoI sites of pGBamdS6. The resulting plasmid was designated pGBamdS7 (Figure 40).

[0185] To destroy the HindIII site approximately 66 bp upstream the startcodon from the chymosin gene, pGBamdS7 was partially digested with HindIII and treated with the Klenow fragment of E.coli DNA polymerase I to generate blunt ends, subsequently ligated and transferred to E.coli for molecular cloning. The derived plasmid was designated pGBamdS8 and contains a LAC4 promoter fragment with a destroyed HindIII site.

Example 9

Expression of the amdS cDNA from the LAC4 promoter

in the yeast K.lactis

[0186] The expression vector, pGBamdS1 contains, apart from the amdS cDNA a second selection marker which confers resistance to the antibiotic G418. This allows to first select for transformants using G418 resistance which is a well established procedure (Sreekrishna et al. (1984) Gene 28, 73-81). The transformants obtained this way can subsequently be used to verify expression of the amdS cDNA and to optimize conditions for selection of the amdS⁺ phenotype in K.lactis. Once these conditions have been established, direct selection for amdS⁺ transformants can be performed, e.g. using expression cassettes without additional selection markers.

[0187] pGBamdS1 (Figure 34) was linearized in the LAC4 promoter by SstII digestion. The pTZ19R sequences were removed by fractionation in and purification from agarose gels. 15 µg's of this DNA fragment were used to transform to the K.lactis strains CBS 2360 and CBS 683 as described by Ito H. et al. (1983) J. Bacteriol. 153, 163-168 with the modifications described under Experimental. The transformation plates were incubated at 30°C for 3 days. G418-resistant transformants were obtained with both strains. Several independent transformants of both strains as well as the wild type strains were subsequently streaked onto plates containing different solid media (see Table 1). YEPD and YEPD/G418 have been described in Experimental. YCB, YCB/NH₄ and YCB/acetamide have been described in example 7 as media I, II and III, respectively. YNB-lac/NH₄ and YNB-lac/acetamide contain 0.17 % (w/v) Yeast Nitrogen Base w/o Amino Acids and Ammonium Sulfate (Difco) supplemented with 1 % (w/v) lactose, 30 mM Sodium Phosphate buffer at pH 7.0 and either 0.1 % (w/v) NH₄(SO₄)₂ or 5 mM acetamide, respectively.

[0188] The amdS⁺ phenotype of the CBS 683/pGBamdS1 transformants was obvious on YCB/acetamide (see Table 1). However, the CBS 2360 transformants containing the same expression vector did not show any growth on YCB/acetamide. We reasoned that this might be due to the lack of induction of the LAC4 promoter driving the amdS cDNA in the absence of lactose or galactose as carbon-source dependent differences in the regulation of the LAC4 promoter between different K.lactis strains have been described (Breunig (1989) Mol. Gen. Genet. 216, 422-427). Table 1 shows that this is indeed the case, on medium containing lactose as sole carbon-source and acetamide as sole nitrogen source the CBS 2360 transformants were able to grow. We can therefore conclude that, depending on the carbon-source used, these transformants sufficiently express the A.nidulans amdS cDNA in order to sustain growth of the yeast K.lactis on acetamide as sole nitrogen-source.

[0189] Southern analyses were performed to verify whether integration in the LAC4 promoter had occurred. High molecular weight DNA of several CBS 2360 and CBS 683 transformants was isolated, digested with HindIII and subsequently fractionated by electrophoresis on a 0.7% agarose gel. After transfer to nitrocellulose, hybridization was performed according to standard procedures. As probe a ³²P-labelled approximately 1.5 kb SacII/HindIII LAC4 promoter fragment isolated from plasmid pGBHSA20 (Figure 34) was used. We identified CBS 683 and CBS 2360 transformants containing a single pGBamdS1 expression cassette integrated in the LAC4 locus, one example of each is shown in figure 41 and is designated KAM-1 and KAM-2, respectively. Single copy integration of pGBamdS1 in the LAC4 promoter produces two new HindIII fragments of 4.2 and 8.6 kb, both of which are present in transformants KAM-1 and KAM-2. Since CBS 683 contains two LAC4 loci and pGBamdS1 has integrated in only one of them in KAM-1, the digest of KAM-1 also shows the 5.6 kb HindIII fragment derived from the second undisturbed LAC4 locus.

Table 1

Growth of <u>K.lactis</u> CBS 683 and CBS 2360 wild type and pGBamdS1 transformants on solid media containing different nitrogen- and/or carbon-sources.				
strain	CBS 683		CBS 2360	
transforming DNA	none	pGBamdS1	none	pGBamdS1
YEPD	+	+	+	+
YEPD-G418	-	+	-	+
YCB	-	-	-	-

Table 1 (continued)

Growth of <i>K.lactis</i> CBS 683 and CBS 2360 wild type and pGBamdS1 transformants on solid media containing different nitrogen- and/or carbon-sources.				
strain	CBS 683		CBS 2360	
YCB / NH ₄	+	+	+	+
YCB / acetamide	-	+	-	-
YNB-lac / NH ₄	+	+	+	+
YNB-lac / acetamide	-	+	-	+

Example 1015 Direct selection of *K.lactis* CBS 683 and CBS 2360 transformants using acetamide as sole nitrogen-source

[0190] SstII linearized pGBamdS1 (15 µg) was transformed into *K.lactis* CBS 683 and CBS 2360 using the transformation procedure as described by Ito H. et al. (1983). *J. Bact.* 153, 163-168.) with the following modifications:

- 20 - *K.lactis* cultures were harvested for transformation at OD₆₁₀ = 0.5 - 1.0.
- After the 5 minutes heatshock of the DNA-cell suspension, the phenotypic expression prior to plating was performed for 150-180 minutes at 30°C in volumes of 1 ml. Different media were used for both strains. For CBS 683 a YEPD/ YNB solution (1*YNB (Yeast Nitrogen Base, Difco), 1% bacto-peptone, 1% yeast extract and 2% glucose) or YNB-glu (1*YNB (Yeast Nitrogen Base w/o Amino Acids and Ammonium Sulphate, Difco) supplemented with 1 % (w/v) glucose and 30mM Sodium Phosphate buffer at pH 7.0) were used. After this incubation the cells were centrifuged at 2000 g at room temperature for 5 minutes and subsequently plated on YCB/acetamide (see example 7). For CBS 2360, YNB-lac (1*YNB (Yeast Nitrogen Base w/o Amino Acids and Ammonium Sulphate, Difco) supplemented with 1 % (w/v) lactose and 30mM Sodium Phosphate buffer at pH 7.0) was used. After this incubation the cells were centrifuged at 2000 g at room temperature for 5 minutes and subsequently plated on YNB-lac/acetamide (see example 9).

Growth was performed at 30°C for 3 days. amdS⁺ transformants were obtained for both strains. The transformation frequencies found were comparable to that found when using the G418 selection. The correct identity of the transformants was confirmed by subsequent plating on YEPD-plates containing G418 and by Southern analysis.

35 [0191] pGBamdS3 (Figure 36), in which the amdS cDNA is driven from the KIEF1 promoter, was linearized in the LAC4 terminator by digestion with XhoI and 15 µg of the gel-isolated fragment was subsequently transformed into the *K.lactis* strain CBS 683 using direct selection on YCB/acetamide plates as described above for the transformation of pGBamdS1 into CBS 683. Some of the transformants obtained were analyzed by Southern blotting. High molecular weight DNA was isolated, digested with BamHI and subsequently separated by electrophoresis on a 0.7% agarose gel. Following transfer to nitrocellulose, hybridization was performed according to standard procedures. As probe the ³²P-labelled 1.2 kb SphI/HindIII LAC4 terminator fragment isolated from plasmid pTY75LAC4 (described in Example 8) was used. The results of several CBS 683 transformants, containing the pGBamdS3 plasmid and from several transformants containing the pGBamdS5 plasmid are shown in figure 42A and 42B respectively. The reference strain CBS 683 is shown in figure 42B. In the CBS 683 transformants an additional 6.8 kb sized hybridizing fragment is present besides the 3.7 kb hybridizing fragment of the intact LAC4 terminator. This implicates a correct integration of the plasmids into the LAC4 terminator region.

45 [0192] In all of these transformants pGBamdS3 was integrated in one or more copies into the LAC4 terminator (the intensity of the 6.8 kb hybridizing fragment is an indication for the number of integrated copies of the vector). We conclude that also the constitutive KIEF1 promoter can drive the amdS cDNA for use as selection marker. Similar results were obtained with pGBamdS5 (Figure 37), in which the amdS cDNA is driven from the S.cerevisiae ADH1 promoter.

Example 1155 Transformation of *S.cerevisiae* with pGBamdS5 by direct selection on acetamide

[0193] In this example we have tested whether the amdS cDNA can also be used as selection marker in other yeasts,

e.g. *S.cerevisiae*. We have first established the *amdS*⁻ phenotype of *S.cerevisiae* strain D237-10B and its ability to use ammonium as sole nitrogen-source, using the same media and procedures as we have described for *K.lactis* in example 7. As observed in the case of *K.lactis*, full grown *S.cerevisiae* colonies were only observed on the plates containing ammonium as nitrogen source. In plates without nitrogen-source or with acetamide as sole nitrogen-source no growth or, occasionally a slight background growth was observed. Plasmid pGBamdS5 was linearized in the *ADH1* promoter by partial digestion with *SphI*. The *S.cerevisiae* strain D273-10B (ATCC 25657) was transformed with 15 µg of gel-isolated linearized pGBamdS5 fragment, using transformation procedures as described in example 10 for the transformation of pGBamdS1 to *K.lactis* CBS 683. After transformation the cells were plated onto YCB/acetamide plates (see Example 9) and allowed to grow at 30°C for 3 days. Several *amdS*⁺ transformants were obtained with in this transformation. Subsequent Southern analysis of some of the *amdS* transformants confirmed that the *amdS* cDNA was stably integrated into the *S.cerevisiae* genome.

[0194] High molecular weight DNA was isolated and digested with *BamHI*, subsequently separated by electrophoresis on a 0.7% agarose gel and blotted onto nitrocellulose. As probe the ³²P labelled 750 bp *EcoRV* *amdS* fragment was used isolated from pGBamdS1. The results of several D273-10B/pGBamdS5 transformants as well as the reference strain D273-10B (ATCC 25657) are shown in figure 43. Two hybridizing fragments are present in the D273-10B transformants respectively, a 6.6 kb fragment that represents the multicopy fragment and a hybridizing fragment of unknown size that represents the flanking. The reference strain D273-10B (ATCC 25657) as expected does not show any hybridizing fragment.

Example 12

Removal of the *amdS*-marker from *K.lactis* and *S.cerevisiae* *amdS*⁺ transformants using fluoracetamide counter-selection

[0195] In the above described examples the *amdS* containing expression cassettes are integrated by a single cross-over homologous recombination in the *K.lactis* and *S.cerevisiae* genomes. This means that the *amdS* cDNA is flanked by direct repeats in the genomes of these *amdS* yeast transformants. Consequently, the *amdS* cDNA will be deleted in a small fraction of the transformant population by intra-chromosomal mitotic recombination events occurring at low frequency between the direct repeats flanking the cDNA. It should be possible to select for these events using media containing fluoracetamide, a compound which is toxic for *amdS*⁺ cells but not for *amdS*⁻ cells as has been shown for *A.nidulans* by Hynes and Pateman ((1970) Mol. Gen. Genet. 108, 107-116). In *amdS*⁺ cells fluoracetamide is converted into ammonium and fluoracetate, the latter being toxic when activated by the enzyme acetyl-CoA-synthetase. Prerequisites for the fluoracetamide counter-selection to also work on *amdS*⁺ yeasts are therefore 1) fluoracetamide should not be toxic for *amdS*⁻ yeasts, 2) the yeast cellwall and plasmamembrane should be permeable to fluoracetamide and 3) the enzyme acetyl-CoA-synthetase should be active. To test this we have used a *K.lactis* CBS 683 transformant containing a single copy of pGBamdS1 integrated in the *LAC4* promoter, designated KAM-1 and a *S.cerevisiae* D273-10B transformant containing a single copy of pGBamdS5 integrated in the *ADH1* promoter, designated SAM-1. Both KAM-1 and SAM-1 were subjected to at least 3 rounds of genetic purification on selective medium (YCB/acetamide) to exclude contamination with the *amdS*⁻ parental strains. KAM-1 and SAM-1 were each plated at a density of approximately 10³ CFU per plate onto YCB/NH₄ supplemented with 10 mM fluoracetamide. For both KAM-1 and SAM-1, 5 to 20 fluoracetamide resistant colonies appeared after 3 to 6 days at 30°C. Southern analysis on chromosomal DNA from several independent KAM-1 and SAM-1 derived *amdS*⁻ colonies confirmed that the *amdS* cDNA was correctly removed from the *K.lactis* and *S.cerevisiae* genomes by homologous recombination between the flanking direct repeats (Figure 41). In fact, in one of the KAM-1 *amdS*⁻ recombinants the crossover-point of the recombination was located between a polymorphic *HindIII* site and the *amdS* cDNA. This polymorphic *HindIII* is present 92 bp upstream of the *LAC4* reading frame in the *LAC4* promoter of pGBamdS1, however, this site is not present in the CBS 683 *LAC4* promoter. The recombination event has left the *HindIII* site in the genome of this particular KAM-1 recombinant which could otherwise not be discriminated from the parent strain CBS 683 (see the extra 4.2 kb fragment in figure 41, lane 6). This KAM-1 recombinant therefore excludes the possibility that we would have isolated CBS 683 contaminants instead of KAM-1 *amdS*⁻ recombinants. We conclude from the above that the *amdS* cDNA can be removed from yeast genomes when flanked by direct repeats using fluoracetamide counter-selection. In the present example the *amdS*⁻ *K.lactis* and *S.cerevisiae* recombinants occur at a frequency of about 0.1 %.

[0196] We have noted that for some yeast strains efficient counter-selection on fluoracetamide cannot be performed on YCB/NH₄, probably due to strong carbon-catabolite repression of the acetyl-CoA-synthetase. In those instances we have successfully used YNB-galactose/NH₄ (this medium is identical to YNB-lac/NH₄ described in example 9 but contains 1 % galactose instead of 1 % lactose) supplemented with 10 mM fluoracetamide for counter-selection.

Example 13Marker gene free deletion of a K.lactis gene using the amdS-marker

5 **[0197]** A frequently used technique for the manipulation of yeast genomes is "one-step gene disruption", a method which allows to disrupt (or modify) genes in a single transformation step (Rothstein et al. (1983) Methods Enzymol. 101, 202-211). In this method a transforming plasmid with a copy of a target gene disrupted by a yeast selectable marker integrates into the yeast genome via a double cross-over homologous recombination, resulting in the replacement of the wild-type target gene by the disrupted copy. Combination of "one-step gene disruption" and the fluoracetamide counter-selection of amdS⁺-yeast-transformants as we have described in Example 12, should enable the deletion of genes from yeast genomes without leaving selectable markers. In this example we have used this combination to delete the LAC4 gene from the K.lactis CBS 2360 genome. For one-step gene transplacement of the K.lactis LAC4 gene pGBamdS6 (Figure 38) was constructed, which contains the amdS expression-cassette flanked by LAC4 promoter and terminator sequences. An additional LAC4 terminator fragment is present directly upstream of the amdS expression-cassette such that the amdS expression-cassette is flanked by direct repeats which will allow the excision of the amdS sequences from the K.lactis genome by intrachromosomal recombination between these direct repeats. Plasmid pGBamdS6 was digested with SpeI and HindIII and a 6.6 kb DNA fragment was isolated after gel electrophoresis. This SpeI-HindIII fragment, containing the gene replacement vector, was used to transform K.lactis CBS 2360 using transformation procedures described in Example 10. amdS⁺ transformants were plated onto on YEPD plates containing 0.008 % X-gal (5-bromo-4-chloro-3-indolyl β-D-galactopyranoside) in order to screen for transformants with a transplaced LAC4 gene.

15 **[0198]** The amdS⁺ transformants were analyzed on Southern blot. High molecular weight DNA was isolated, digested with HindIII, subsequently separated by electrophoresis on a 0.7% agarose gel and blotted onto nitrocellulose. As probe a ³²P-labelled 600 bp XbaI LAC4 terminator fragment isolated from plasmid pPTLAC4 (described in example 8) was used. The results of an amdS⁺ CBS 2360 transformant with a transplaced LAC4 gene as well as the reference strain CBS 2360 are shown in figure 44. In case of the amdS⁺ CBS 2360 transformant, a 7.4 kb hybridizing fragment is present that implicates a correctly transplaced LAC4 gene. The reference strain CBS 2360 shows a 2.0 kb hybridizing fragment that represents the intact LAC4 locus.

25 **[0199]** Subsequent fluoracetamide counter-selection of these amdS⁺ transformants as described in Example 12, yielded recombinants with an amdS⁻ phenotype. Southern analysis was performed on the chromosomal DNA of the amdS⁻ recombinants. High molecular weight DNA was isolated, digested with HindIII, subsequently separated on a 0.7% agarose gel and blotted onto nitrocellulose. The same ³²P-labelled probed as described above was used. The results of the amdS⁻ CBS 2360 recombinants are shown in figure 44. In case of the amdS⁻ recombinants, a 5.4 kb hybridizing fragment is present, which confirmed the absence of the LAC4 gene as well as the correct removal of the amdS marker from the yeast genome. The absence of the amdS marker from these K.lactis LAC4⁻ strains offers the possibility to reuse the amdS marker for additional deletions and/or modifications of genes.

Example 14Marker gene free insertion of a gene into the K.lactis genome using the amdS marker

40 **[0200]** For the marker gene free insertion of genes into the yeast genome we have used the chymosin cDNA as a model-gene. In this example we have inserted the chymosin cDNA at the K.lactis LAC4 locus while replacing the LAC4 gene and without leaving a selection marker. The principle of marker-free gene insertion is the same as that for marker-free deletion of genes as described in example 13 except that in this case the transplacement vector pGBamdS8 contains a gene of interest, the chymosin cDNA (Figure 40). Plasmid pGBamdS8 was digested with SpeI and HindIII and the 8.0 kb DNA fragment was gel-isolated. 10 µg of this fragment was transformed to K.lactis CBS 2360 as described in Example 10. amdS⁺ transformants with a transplaced LAC4 gene and chymosin activity were obtained. Chymosin activity was measured as described (van den Berg et al. (1990) Bio/technology 8, 135-139). By subsequent counter-selection of these transformants on fluoracetamide as described in example 12 recombinants were isolated with an amdS⁻ phenotype but which still produced chymosin. Southern analysis of the chromosomal DNA of the amdS⁻, Chymosin⁺ recombinants confirmed the replacement of the LAC4 gene by the chymosin cDNA as well as the correct removal of the amdS marker from the K.lactis genome. The amdS⁻/chymosin⁺ phenotype of these recombinants was also confirmed by lack of growth on YCB/acetamide plates and by the presence of chymosin activity (see above). The amdS⁻ phenotype of these recombinants allows further manipulation of these strains using the amdS marker, e.g. integration of additional copies of the chymosin expression-cassette and/or deletion of K.lactis genes as described in example 13.

(Figure 50) was isolated.

Construction of pGBamdS41

5 [0206] For the expression of the *A.nidulans* *amdS* cDNA in *E.coli* we have used pTZ18R/N, a derivative of pTZ18R which is described in the European Patent Application 0 340 878 A1 pTZ18R/N differs from pTZ18R in that an *NdeI* site was created at the ATG start-codon of the *lacZ* reading frame in pTZ18R using *in vitro* site directed mutagenesis. pGBamdS21 was digested with *NdeI* and *HindIII* and the gel-isolated fragment containing the *amdS* cDNA was ligated into pTZ18R/N digested with *NdeI* and *HindIII*. This ligation mixture was used to transform *E.coli* JM109 and from one of the transformants pGBamdS41 (Figure 51) was isolated.

Example 17

Transformation of Bacilli using the *amdS* gene as selection marker

15 [0207] In order to delete the *E.coli* sequences from pGBamdS22 and to place the "hpa2"-promoter immediately upstream of the *amdS* cDNA, pGBamdS22 was digested with *NdeI*, recircularized by ligation and used to transform *B.subtilis* BS-154 (CBS 363.94). Transformants were selected on acetamide minimal plates and checked for neomycin resistance. From one of these transformants expression-vector pGBamdS23 (Figure 49) was isolated and characterized by restriction enzyme analysis. These results show that 1) the *A.nidulans* *amdS* cDNA under the control of a *Bacillus* promoter sequence is expressed well and 2) that the *amdS* gene can be used as a selection marker in the transformation of *Bacilli*.

20 [0208] *B.licheniformis* T5 (CBS 470.83) was transformed with vector pGBamdS25. Transformation was performed as described in Experimental and *amdS*⁺ transformants were obtained by direct selection on modified protoplast regeneration plates supplemented with 20 mM acetamide as sole nitrogen source (described in Experimental). The presence of pGBamdS25 in the transformants was confirmed by their neomycin resistance phenotype as well as the fact that the plasmid could be reisolated from the transformants. One of these transformants designated BAA-103 was used to achieve integration of plasmid pGBamdS25 into the *B.licheniformis* genome targeted at the amylase locus. Plasmid integration was performed by growing transformants at 50°C on minimal medium agar containing acetamide as sole nitrogen source. Several colonies were transferred repeatedly (2 to 3 times) to fresh plates followed by incubation at 50°C. Isolated colonies were tested for their ability to grow on acetamide as sole nitrogen source and for resistance to neomycin at 1 µg/ml. The absence of autonomously replicating plasmid DNA was established by re-transformation of DNA isolated from the integrants to the host strain. No neomycin resistant colonies could be obtained.

30 [0209] This result is a clear evidence that the *amdS* gene is a suitable marker to select *Bacillus* species containing a single *amdS* gene copy.

Example 18

Transformation of *E.coli* using the *amdS* gene as selection marker

40 [0210] *E.coli* JM109 was transformed with the vector pGBamdS41 using standard procedures. Selections were performed on either M9 plates supplemented with 0.02 µg/ml thiamine and 50 µg/ml ampicillin or M9 plates without ammonium but supplemented with 20 mM acetamide, 0.02 µg/ml thiamine and 0.05 mM IPTG. Several *amdS*⁺/ampicillin resistant transformants were obtained from which pGBamdS41 could be reisolated. The transformation frequencies using selection on ampicillin or acetamide were comparable. This demonstrates that the *A.nidulans* *amdS* gene is functional as selection marker for the transformation of Gram-negative bacteria as well.

Example 19

Fluoracetamide counter-selection of *amdS*⁺ bacterial transformants

50 [0211] Counter-selection of bacterial *amdS*⁺ transformants using fluoracetamide requires the activity of the enzyme acetylCoA synthetase for the conversion of fluoracetate to fluoracetyl-CoA. To avoid catabolite repression of acetylCoA synthase, as has been observed in *E.coli* (Brown et al., 1977), bacterial *amdS*⁺ transformants or single copy integrants were grown on defined media containing NH₄Cl as nitrogen source and acetate as carbon and energy source.

55 [0212] Many organisms including *B. subtilis* (Freese, E. and Fortnagel, P. (1969) J. Bacteriol 90, 745-756) lack a functional glyoxylate shunt and therefore metabolize acetate only when the medium is supplemented with a source of TCA cycle intermediates, such as glutamate or succinate. *Bacillus amdS*⁺ strains were grown on TSS medium with

EP 1 321 523 A2

0.01% glutamate and 50 mM acetate as described by Grundy, F.J. et.al. (1993) Molecular Microbiology 10, 259-271. To this medium solidified with agar, fluoracetamide was added in concentrations ranging from 1 to 50 mM. B.subtilis BAA-101 or B.licheniformis BAA-103 (single copy integrant) were plated at a density of 10^2 cells per plate. At a certain fluoracetamide concentration only a few colonies appeared. The absence of pGBamdS25 in these colonies was demonstrated by plasmid and chromosomal DNA analysis, sensitivity towards neomycin, and inability to grow on acetamide as sole nitrogen-source. Counter-selection of BAA-103 in some cases led to the loss of the amylase gene as indicated by activity assays and Southern blots. This shows that fluoracetamide counter-selection can be used to select amdS⁻ cells from a population containing a majority of amdS⁺ Bacillus cells and the simultaneous deletion of a specific target gene.

10 [0213] Similarly we have used minimal medium #132 as described by Vanderwinkel E. and De Vlieghe M, European J. Biochem, 5 (1968) 81-90 supplemented with fluoracetamide in concentrations ranging from 1 to 50 mM and 0.05 mM IPTG to select amdS⁻ E.coli JM109 cells from a population of pGBamdS41 transformants. Cells were plated at a density of 10^2 cells per plate. At a certain fluoracetamide concentration only a few colonies appeared. The absence of pGBamdS41 from the fluoracetamide selected colonies was confirmed by isolation of DNA, sensitivity toward ampicillin and inability to grow on acetamide as sole nitrogen-source. This demonstrates that the fluoracetamide counter-selection can be used to select amdS⁻ cells from a population containing a majority of amdS⁺ E.coli cells.

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Gist-brocades B.V.
- (B) STREET: Wateringseweg 1
- (C) CITY: Delft
- (E) COUNTRY: The Netherlands
- (F) POSTAL CODE (ZIP): 2611 XT

(ii) TITLE OF INVENTION: Selection Marker Gene Free Recombinant
Strains: a Method for Obtaining Them and the Use of These
Strains

(iii) NUMBER OF SEQUENCES: 37

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (B) CLONE: AB3100

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CTAATCTAGA ATGCCTCAAT CCTGAA

26

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:

- (B) CLONE: AB3101

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GACAGTCGAC AGCTATGGAG TCACCACA

28

(2) INFORMATION FOR SEQ ID NO: 3:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 10 (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (vii) IMMEDIATE SOURCE:
15 (B) CLONE: TN0001

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

20 TCGATTAAGT AGTTAA 16

(2) INFORMATION FOR SEQ ID NO: 4:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 30 (iii) HYPOTHETICAL: NO
- (vii) IMMEDIATE SOURCE:
(B) CLONE: AB2154

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

35 AACCATAGGG TCGACTAGAC AATCAATCCA TTTCG 35

(2) INFORMATION FOR SEQ ID NO: 5:

- 40 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 45 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: YES
- 50 (vii) IMMEDIATE SOURCE:
(B) CLONE: AB2155

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

55 GCTATTCGAA AGCTTATTCA TCCGGAGATC CTCAT 35

(2) INFORMATION FOR SEQ ID NO: 9:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 10 (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: YES
- (vii) IMMEDIATE SOURCE:
15 (B) CLONE: AB2994

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

20 CTGATAGAAT TCAGATCTGC AGCGGAGGCC TCTGTG 36

(2) INFORMATION FOR SEQ ID NO: 10:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 30 (iii) ANTI-SENSE: NO
- (vii) IMMEDIATE SOURCE:
(B) CLONE: AB3657

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

35 AGCTTGACGT CTACGTATTA ATGCGGCCGC T 31

(2) INFORMATION FOR SEQ ID NO: 11:

- 40 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
45 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: YES
- 50 (vii) IMMEDIATE SOURCE:
(B) CLONE: AB3658

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

55 TCGAAGCGGC CGCATTATA CGTAGACGTC A 31

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3779

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

AATTGGGGCC CATTAACTCG AGC

23

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3780

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

AATTGCTCGA GTTAATGGGC CCC

23

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3448

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GTGCGAGGTA CCACAATCAA TCCATTTCGC

30

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3449

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

ATGGTTC AAG AACTCGGTAG CCTTTCCTT GATTCT

36

(2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3450

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

AGAATCAAGG AAAAGGCTAC CGAGTCTTG AACCAT

36

(2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3520

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

ATCAATCAGA AGCTTCTCT CGAGACGGGC ATCGGAGTCC CG

42

(2) INFORMATION FOR SEQ ID NO: 18:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 10 (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (vii) IMMEDIATE SOURCE:
 15 (B) CLONE: AB3781

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

AATTGGGGCC CAGCGTCC

18

20

(2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 25 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 30 (iii) ANTI-SENSE: YES
- (vii) IMMEDIATE SOURCE:
 (B) CLONE: AB3782

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

AATTGGACGC TGGGCCCC

18

35

(2) INFORMATION FOR SEQ ID NO: 20:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- 45 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- 50 (vii) IMMEDIATE SOURCE:
 (B) CLONE: AB3746

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

55

TGACCAATAA AGCTTCTCGA GTAGCAAGAA GACCCAGTCA ATC

43

(2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 47 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3747

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

CTACAAACGG CCACGCTGGA GATCCGCCGG CGTTCGAAAT AACCAGT

(2) INFORMATION FOR SEQ ID NO: 22:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB4234

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GAAGACCCAG TCAAGCTTGC ATGAGC

(2) INFORMATION FOR SEQ ID NO: 23:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB4235

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

TGACCAATTA AGCTTGCGGC CGCTCGAGGT CGCACC GGCA AAC

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (B) CLONE: AB4236

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

TGACCAATAA AGCTTAGATC TGGGGGTGAT TGGGCGGAGT GTTTTGCTTA GACAATCAAT 60
CCATTTCGC 69

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:

- (B) CLONE: AB4233

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

TGACCAATAG ATCTAAGCTT GACTGGGTCT TCTTGC 36

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (B) CLONE: AB3514

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

CTGCCGAATTC GTCGACATGC CTCATCCTG GG 32

5

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

15 (iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:

(B) CLONE: AB3515

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

GGCAGTCTAG AGTCGACCTA TGGAGTCACC ACATTTC 37

25

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

35

(vii) IMMEDIATE SOURCE:

(B) CLONE: AB3701

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

40 CTGCCGAATTC GTCGACACTA GTGGTACCAT TATAGCCATA GGACAGCAAG 50

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 70 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: DNA (genomic)

50

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:

(B) CLONE: AB3700

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

GCTCTAGAGC GCGCTTATCA GCTTCCAGTT CTTCCCAGGA TTGAGGCATT TTTAATGTTA 60
 5 CTTCTCTTGC 70

(2) INFORMATION FOR SEQ ID NO: 30:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 50 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 15 (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (vii) IMMEDIATE SOURCE:
 20 (B) CLONE: AB3702

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

CTGCGAATTC GTCGACACTA GTGGTACCAT CCTTTTGTG TTTCCGGGTG 50

(2) INFORMATION FOR SEQ ID NO: 31:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 70 base pairs
 (B) TYPE: nucleic acid
 30 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 35 (iii) ANTI-SENSE: YES
- (vii) IMMEDIATE SOURCE:
 (B) CLONE: AB3703

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

GCTCTAGAGC GCGCTTATCA GCGGCCAGTT CTTCCCAGGA TTGAGGCATT GTATATGAGA 60
 TAGTTGATTG 70

(2) INFORMATION FOR SEQ ID NO: 32:

- 45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 55 base pairs
 (B) TYPE: nucleic acid
 50 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 55

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

(B) CLONE: AB3704

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

GCTCTAGAAG TCGACACTAG TCTGCTACGT ACTCGAGAAT TTATACTTAG ATAAG

(2) INFORMATION FOR SEQ ID NO: 33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:

(B) CLONE: AB3705

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

TGCTCTAGAT CTCAGCCAC AATTC

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

(B) CLONE: AB3965

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

CTGCTACGTA ATGTTTTTCAT TGCTGTTTTTA C

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3966

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

CCGCCCCAGTC TCGAGTCAGA TGGCTTTGGC CAGCCCC

(2) INFORMATION FOR SEQ ID NO: 36:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3825

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

CGCGCTTATC AGCGGCCAGT TCTTCCCAGG ATTGAGGCAT ATGT

(2) INFORMATION FOR SEQ ID NO: 37:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(vii) IMMEDIATE SOURCE:
(B) CLONE: AB3826

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

CTAGACATAT GCCTCAATCC TGGGAAGAAC TGGCCGCTGA TAAG

Claims

1. A selection marker free recombinant filamentous fungus, which contains a recombinant vector comprising a desired DNA fragment.
2. The selection marker free recombinant filamentous fungus according to claim 1, **characterized in that** the desired

DNA fragment is integrated into the genome of said filamentous fungus through site specific homologous recombination.

- 5 3. The selection marker free recombinant filamentous fungus according to any one of claim 1 to 2, **characterized in that** the desired DNA fragment contains a genetic element selected from the group consisting of: a gene, a cDNA, a promoter, a terminator, a regulatory element, an intron, a recognition sequence for a DNA-binding protein, a translation-initiation site, a restriction site and combinations thereof.
- 10 4. The selection marker free recombinant filamentous fungus according to any one of claims 1 to 3, **characterized in that** the desired DNA fragment contains a sequence encoding a chymosin, a phytase, a xylanase, a lipase, an amylase, a protease or a β -galactosidase.
- 15 5. The selection marker free recombinant filamentous fungus according to any one of claim 1 to 4, **characterized in that** the desired DNA fragment is foreign to said filamentous fungus.
6. A selection marker free recombinant filamentous fungus preferably according to any one of claims 1 to 5, **characterized in that** it comprises a desired deletion in its genome.
- 20 7. A selection marker free recombinant filamentous fungal strain according to any one of claim 1 to 6, wherein the filamentous fungus is an *Aspergillus*, *Trichoderma*, or *Penicillium* species.

25

30

35

40

45

50

55




Fig. 1




Fig. 2




Fig. 3a




Fig. 3b




Fig. 4




Fig. 5




Fig. 6




Fig. 7a




Fig. 3b




Fig. 4




Fig. 5




Fig. 6




Fig. 7a




Fig. 7b




Fig. 8a




Fig. 8b




Fig. 9




Fig. 10a




Fig. 10b




Fig. 11




Fig. 12a




Fig. 12b




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17a




Fig. 17b




Fig. 18




Fig. 19




Fig. 20

5' AATTGGGGCCCAGCGTCC 3'
3' CCCCgggTcgcAGGTAA 5'




Fig. 21




Fig. 22




Fig. 23a




Fig. 23b




Fig. 24




Fig. 25




Fig. 26a




Fig. 26b




Fig. 27




Fig. 28




Fig. 29




Fig. 30a




Fig. 30b




Fig. 31




Fig. 32a




Fig. 32b




Fig. 33




Fig. 34




Fig. 35




Fig. 36




Fig. 37




Fig. 38




Fig. 39




Fig. 40




Fig. 41




Fig. 42




Fig. 43




Fig. 44




Fig. 45




Fig. 46




Fig. 47




Fig. 48




Fig. 49




Fig. 50




Fig. 51